



ATLANTIC HEALTH DATABASE

APPENDIX C

DEATH & DISEASE

LIST OF SYMBOLS

—	Nil or zero.
--	Number is too small to be expressed.
·	Data are not available for any reference period. This symbol is used when the figure has never been published for any reference period. The symbol implies that the underlying time series does not exist, but theoretically could exist.
..	Data are not available for a specific reference period. This symbol is used when the time series exists but the figure is not published for a specific reference period.
...	Not applicable. This symbol is used when an entry for the table cell is not logically possible. For example, there could not be any data for "legal drivers under 15 years of age" or "women with prostate cancer." In such cases, no time series will exist.
×	Used to rank a region relative to a province or Canada. In tables showing percentages, × denotes a lower percentage than the province or nation. In tables showing rates of occurrence, × denotes a less favourable rate.
✓	Used to rank a region relative to a province or Canada. In tables showing percentages, ✓ denotes a higher percentage than the province or nation. In tables showing rates of occurrence, ✓ denotes a more favourable rate.
-	In tables comparing a region relative to a province or Canada, “-” denotes that there is no data available for comparison
^E	Data with a coefficient of variation (CV) from 16.6% to 33.3% are identified by ^E and should be interpreted with caution.
F	Data with a coefficient of variation (CV) greater than 33.3% were suppressed due to extreme sampling variability. When a figure is "too unreliable to be published," the data point is suppressed and the symbol F appears in the data cell.
s	Used in tables ranking regions relative to a province or Canada. In tables showing percentages, s denotes that the percentage is equal to province or nation. In tables showing rates of occurrence, s denotes an equal rate.
x	Figures have been suppressed to meet the confidentiality provisions of the <i>Statistics Act</i> .

LIST OF TABLES

Table 113. Age standardized total mortality rate (ICD-9 001-799, E800-E999), Canada, Atlantic Provinces, and Health Districts, 1996	10
Table 114. Total mortality (ICD-9 001-799, E800-E999), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996.....	11
Table 115. Total mortality (ICD-9 001-799, E800-E999), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	12
Table 116. Total mortality (ICD-9 001-799, E800-E999), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	13
Table 117. Age standardized circulatory disease death rate (ICD-9 390-459), Canada, Atlantic Provinces, and Health Districts, 1996.....	14
Table 118. All circulatory disease deaths (ICD-9 390-459), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996.....	15
Table 119. All circulatory disease deaths (ICD-9 390-459), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	16
Table 120. All circulatory disease deaths (ICD-9 390-459), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	17
Table 121. Ischaemic heart disease deaths (ICD-9 410-414), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996.....	18
Table 122. Ischaemic heart disease deaths (ICD-9 410-414), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	19
Table 123. Ischaemic heart disease deaths (ICD-9 410-414), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	20
Table 124. Cerebrovascular disease deaths (ICD-9 430-438), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996.....	21
Table 125. Cerebrovascular disease (ICD-9 430-438), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	22
Table 126. Cerebrovascular disease deaths (ICD-9 430-438), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	23
Table 127. All other circulatory disease deaths (ICD-9 390-409, 415-429, 439-459), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	24
Table 128. All other circulatory disease deaths (ICD-9 390-409, 415-429, 439-459), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	25

Table 129. All other circulatory disease deaths (ICD-9 390-409, 415-429, 439-459), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	26
Table 130. Age standardized unintentional injury death rate (ICD-9 E800-E929, excluding 870-879), Canada, Atlantic Provinces, and Health Districts, 1996.....	27
Table 131. Unintentional injury deaths (ICD-9 E800-E929, excluding 870-879), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	28
Table 132. Unintentional injury deaths (ICD-9 E800-E929, excluding 870-879), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	29
Table 133. Unintentional injury deaths (ICD-9 E800-E929, excluding 870-879), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996.....	30
Table 134. Age standardized AIDS death rate (ICD-9 042-044), Canada, Atlantic Provinces, and Health Districts, 1996	31
Table 135. AIDS deaths (ICD-9 042-044), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996 ..	32
Table 136. AIDS deaths (ICD-9 042-044), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996.....	33
Table 137. AIDS deaths (ICD-9 042-044), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	34
Table 138. Age standardized cancer (malignant neoplasms) death rate (ICD-9 140-208), Canada, Atlantic Provinces, and Health Districts, 1996.	35
Table 139. All cancer deaths (malignant neoplasms) (ICD-9 140-208), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	36
Table 140. All cancer deaths (malignant neoplasms) (ICD-9 140-208), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	37
Table 141. All cancer deaths (malignant neoplasms) (ICD-9 140-208), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	38
Table 142. Lung cancer deaths (ICD-9 162), by sex, crude count and confidence interval, Canada, Atlantic Provinces, and Health Districts, 1996	39
Table 143. Lung cancer deaths (ICD-9 162), by sex, crude rate per 100,000 population and confidence interval, Canada, Atlantic Provinces and Health Districts, 1996.....	40
Table 144. Lung cancer deaths (ICD-9 162), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	41
Table 145. Colorectal cancer deaths (ICD-9 153-154), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996.....	42
Table 146. Colorectal cancer deaths (ICD-9 153-154), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	43

Table 147. Colorectal cancer deaths (ICD-9 153-154), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	44
Table 148. Prostate cancer deaths (ICD-9 185), crude count, crude rate per 100,000 population, age-standardized rate per 100,000 population and confidence intervals, Canada, Atlantic Provinces, and Health Districts, 1996	45
Table 149. Female breast cancer deaths (ICD-9 174), crude count, crude rate per 100,000 population, age-standardized rate per 100,000 population and confidence intervals, Canada, Atlantic Provinces, and Health Districts, 1996	46
Table 150. Age standardized respiratory disease death rate (ICD-9 460-519), Canada, Atlantic Provinces, and Health Districts, 1996.....	47
Table 151. All respiratory disease deaths (ICD-9 460-519), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996.....	48
Table 152. All respiratory disease deaths (ICD-9 460-519), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	49
Table 153. All respiratory disease deaths (ICD-9 460-519), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	50
Table 154. Pneumonia and influenza deaths (ICD-9 480-487), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996.....	51
Table 155. Pneumonia and influenza deaths (ICD-9 480-487), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	52
Table 156. Pneumonia and influenza deaths (ICD-9 480-487), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	53
Table 157. Bronchitis, emphysema and asthma deaths (ICD-9 490-493), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	54
Table 158. Bronchitis, emphysema and asthma deaths (ICD-9 490-493), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	55
Table 159. Bronchitis, emphysema and asthma deaths (ICD-9 490-493), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	56
Table 160. All other respiratory disease deaths (ICD-9 460-519, excluding 480-487, 490-493), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts	57
Table 161. All other respiratory disease deaths (ICD-9 460-519, excluding 480-487, 490-493), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts.....	58
Table 162. All other respiratory disease deaths (ICD-9 460-519, excluding 480-487, 490-493), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	59
Table 163. Deaths due to medically treatable diseases - cervical cancer (ICD-9 180), age-standardized rate per 100,000 population and confidence interval, women aged 5 to 64, Canada, Atlantic Provinces, and Health Districts, 1996.....	60

Table 164. Deaths due to medically treatable diseases - bacterial infections (ICD-9 001-005, 020-041, 320, 382, 383, 390-392, 680-686, 711, 730), age-standardized rate per 100,000 population and confidence interval, population aged 5 to 64, by sex, Canada, Atlantic Provinces, and Health Districts, 1996.....	61
Table 165. Deaths due to medically treatable diseases - pneumonia and unspecified bronchitis (ICD-9 481-486, 490), age-standardized rate per 100,000 population and confidence interval, population aged 5 to 49, by sex, Canada, Atlantic Provinces, and Health Districts, 1996.....	62
Table 166. Deaths due to medically treatable diseases - hypertensive disease (ICD-9 401-405), age-standardized rate per 100,000 population and confidence interval, population aged 5 to 64, by sex, Canada, Atlantic Provinces, and Health Districts, 1996.....	63
Table 167. Thirty day acute myocardial infraction and 30 day stroke mortality 2002	64
Table 168. Arthritis/rheumatism, asthma, diabetes, and high blood pressure, Canada, Atlantic Provinces, and Health Districts, 2000/01	66
Table 169. Arthritis/rheumatism, by sex, household population aged 12 and over, Canada, Atlantic Provinces, and Health Districts, 2000/01	67
Table 170. Arthritis/rheumatism, by sex, household population aged 12 and over, Canada and Atlantic Provinces, 1994/95-1998/99	68
Table 171. Diabetes, by sex, household population aged 12 and over, Canada, Atlantic Provinces, and Health Districts 2000/01	69
Table 172. Diabetes, by sex, household population aged 12 and over, Canada and Atlantic Provinces, 1994/95-1998/99.....	70
Table 173. Asthma, by sex, household population aged 12 and over, Canada, Atlantic Provinces, and Health Districts 2000/01	71
Table 174. Asthma, by sex, household population aged 12 and over, Canada and Atlantic Provinces, 1994/95-1998/99	72
Table 175. High blood pressure, by sex, household population aged 12 and over, Canada, Atlantic Provinces, and Health Districts 2000/01	73
Table 176. High blood pressure, by sex, household population aged 12 and over, Canada and Atlantic Provinces, 1994/95-1998/99	74
Table 177. Age-standardized cancer (malignant neoplasms) incidence (ICD-9 140-208), Canada, Atlantic Provinces, and Health Districts, 1996.....	75
Table 178. All cancer incidence (malignant neoplasms) (ICD-9 140-208), age-standardized rate per 100,000 population, by sex, Canada, Atlantic Provinces, and Health Districts, 1996	76
Table 179. All cancer incidence (malignant neoplasms) (ICD-9 140-208), age-standardized rate per 100,000 population, by sex, Canada and Atlantic Provinces, 1976-2001	77
Table 180. Lung cancer incidence (ICD-9 162), age-standardized rate per 100,000 population, by sex, Canada, Atlantic Provinces, and Health Districts 1996.....	78
Table 181. Lung cancer incidence (ICD-9 162), age-standardized rate per 100,000 population, by sex, Canada and Atlantic Provinces, 1976-2001	79
Table 182. Colorectal cancer incidence (ICD-9 153-154), age-standardized rate per 100,000 population, by sex, Canada, Atlantic Provinces, and Health Districts, 1996.....	80
Table 183. Colorectal cancer incidence (ICD-9 153-154), age-standardized rate per 100,000 population, by sex, Canada and Atlantic Provinces, 1976-2001	81
Table 184. Prostate cancer incidence (ICD-9 185), age-standardized rate per 100,000 population, Canada, Atlantic Provinces, and Health Districts, 1996	82
Table 185. Prostate cancer incidence (ICD-9 185), age-standardized rate per 100,000 population, Canada and Atlantic Provinces, 1976-2001	83

Table 186. Female breast cancer incidence (ICD-9 174), age-standardized rate per 100,000 population, Canada, Atlantic Provinces, and Health Districts, 1996.....	84
Table 187. Female breast cancer incidence (ICD-9 174), age-standardized rate per 100,000 population, Canada and Atlantic Provinces, 1976-2001	85
Table 188. Rate of positive HIV test reports (per 100,000 population), province/territory and year of test, Canada, 1999-2001	86
Table 189. Number of positive HIV test reports by province/territory and year of test, Canada, 1995-2001	86

DEATHS

All data retrieved from:

<http://www4.statcan.ca/survey/invite3/proceed.cgi?loc=http://www.statcan.ca/english/freepub/82-221-XIE/00502/tables.htm>

and

<http://www4.statcan.ca/survey/invite3/proceed.cgi?loc=http://www.statcan.ca/english/freepub/82-221-XIE/00502/tables.htm>

and

http://secure.cihi.ca/hirpt/jsp/En_HIQuery.jsp

Table 113. Age standardized total mortality rate (ICD-9 001-799, E800-E999), Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes		Male		Female	
	Canada	Province	Canada	Province	Canada	Province
Canada						
Newfoundland and Labrador	×		×		×	
St. John's (NF1)	×	✓	×	✓	×	✓
Eastern (NF2)	×	×	×		×	×
Central (NF3)	×	×	×	✓	×	×
Western (NF4)	×	✓	×	✓	×	×
Grenfell (NF5)	×	✓	×	✓	×	×
Labrador (NF6)	×	×	×	×	×	×
Prince Edward Island	×		×		✓	
PEI/Urban	×	×	×	×	×	×
PEI/Rural	✓	✓	×	✓	✓	✓
Nova Scotia	×		×		×	
South-SW (NS1)	×	✓	×	✓	×	✓
Valley (NS2)	✓	✓	✓	✓	✓	✓
Colch-Cumb-E.Hants (NS3)	×	✓	×	✓	×	✓
Pictou-GASHA (NS4)	×	✓	×	✓	×	✓
Cape Breton (NS5)	×	×	×	×	×	×
Capital (NS6)	×	×	×	✓	×	×
New Brunswick	×		×		×	
Moncton (NB1)	✓	✓	✓	✓	✓	✓
Sussex/Saint John (NB2)	×	×	×	×	×	×
Fredericton (NB3)	×	×	×	×	×	×
Edmundston (NB4)	×	×	×	×	×	×
Campbellton (NB5)	×	×	×	×	×	×
Bathurst (NB6)	✓	✓	✓	✓	✓	✓
Miramichi (NB7)	×	×	×	×	✓	✓

× - less favourable total mortality rate than the province or nation

✓ - more favourable total mortality rate than the province or nation

Table 114. Total mortality (ICD-9 001-799, E800-E999), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Crude count	95% CI		Crude count	95% CI		Crude count	95% CI	
		low	high		low	high		low	high
Canada	212,976	212,456	213,497	111,510	111,134	111,887	101,466	101,107	101,825
Newfoundland and Labrador	4,045	3,974	4,117	2,240	2,187	2,294	1,805	1,758	1,853
St. John's (NF1)	1,277	1,237	1,317	670	641	699	607	580	635
Eastern (NF2)	1,043	1,007	1,080	593	566	621	451	427	475
Central (NF3)	876	843	910	492	467	517	384	362	407
Western (NF4)	650	622	680	362	341	384	288	270	308
Grenfell (NF5)	101	90	113	58	50	67	43	36	51
Labrador (NF6)	98	87	109	65	57	75	32	26	39
Prince Edward Island	1,150	1,113	1,189	617	589	645	533	508	560
PEI/Urban	592	565	620	294	275	314	298	279	318
PEI/Rural	558	532	585	323	303	343	236	219	253
Nova Scotia	7,823	7,723	7,923	4,046	3,975	4,118	3,776	3,708	3,846
South-SW (NS1)	1,305	1,265	1,346	691	662	721	614	587	642
Valley (NS2)	688	659	718	358	337	379	331	311	352
Colch-Cumb-E.Hants (NS3)	969	934	1,004	498	473	523	471	447	496
Pictou-GASHA (NS4)	939	905	974	489	465	514	450	426	474
Cape Breton (NS5)	1,427	1,385	1,470	748	718	780	679	650	709
Capital (NS6)	2,494	2,438	2,551	1,262	1,223	1,303	1,232	1,193	1,272
New Brunswick	5,925	5,838	6,012	3,159	3,096	3,222	2,766	2,707	2,826
Moncton (NB1)	1,402	1,360	1,445	753	723	785	649	620	678
Sussex/Saint John (NB2)	1,547	1,503	1,591	773	742	805	773	742	805
Fredericton (NB3)	1,270	1,231	1,311	677	648	706	594	567	622
Edmundston (NB4)	443	419	467	243	226	261	199	184	216
Campbellton (NB5)	271	253	290	147	133	161	124	112	137
Bathurst (NB6)	599	572	627	343	323	365	255	238	274
Miramichi (NB7)	393	371	416	222	206	239	171	157	187

Table 115. Total mortality (ICD-9 001-799, E800-E999), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	717.8	716.0	719.5	759.0	756.4	761.6	677.3	674.9	679.7
Newfoundland and Labrador	721.6	708.8	734.4	801.7	782.8	821.0	641.9	625.0	659.1
St. John's (NF1)	684.4	663.0	706.2	730.7	699.4	763.1	639.6	610.8	669.4
Eastern (NF2)	837.6	808.7	867.1	957.0	913.5	1,002.00	719.5	682.0	758.4
Central (NF3)	773.1	744.1	803.0	861.1	818.2	905.7	683.6	645.1	723.8
Western (NF4)	702.4	671.9	734.0	782.5	737.2	829.9	622.5	582.1	664.8
Grenfell (NF5)	564.2	503.4	630.2	637.2	548.0	736.5	487.8	408.7	577.6
Labrador (NF6)	380.1	338.4	425.4	492.3	427.0	564.7	260.2	212.2	315.8
Prince Edward Island	844.7	816.9	873.1	918.3	877.3	960.6	773.0	735.9	811.4
PEI/Urban	932.1	889.7	976.0	947.2	886.5	1,010.80	917.7	859.2	979.1
PEI/Rural	768.2	732.2	805.5	893.4	838.7	950.7	644.6	598.5	693.2
Nova Scotia	840.0	829.4	850.8	885.7	870.1	901.5	796.0	781.5	810.7
South-SW (NS1)	1,023.4	991.9	1,055.7	1,096.9	1,050.7	1,144.6	951.6	909.1	995.6
Valley (NS2)	824.6	789.7	860.5	867.5	816.9	920.3	782.6	735.2	832.3
Colch-Cumb-E.Hants (NS3)	912.0	879.4	945.5	951.9	904.7	1,000.8	873.4	828.9	919.6
Pictou-GASHA (NS4)	938.7	904.7	973.7	990.5	941.0	1,041.8	888.2	841.9	936.3
Cape Breton (NS5)	1,000.8	971.2	1,030.9	1,079.90	1,036.2	1,125.0	925.9	886.5	966.6
Capital (NS6)	671.6	656.6	686.9	695.0	673.2	717.3	649.3	628.7	670.4
New Brunswick	786.8	775.3	798.4	845.0	828.2	862.1	729.4	713.9	745.2
Moncton (NB1)	766.7	743.9	790.1	832.1	798.4	866.8	702.7	672.0	734.3
Sussex/Saint John (NB2)	869.7	845.0	894.8	887.1	851.7	923.6	852.8	818.8	887.9
Fredericton (NB3)	768.1	744.0	792.6	819.2	784.3	855.3	717.0	684.4	750.8
Edmundston (NB4)	809.1	766.6	853.3	900.1	836.8	966.9	720.2	664.4	779.5
Campbellton (NB5)	822.5	767.6	880.2	902.0	820.9	988.7	745.0	672.5	823.2
Bathurst (NB6)	670.7	640.3	702.1	763.9	718.4	811.4	576.1	536.5	617.9
Miramichi (NB7)	786.9	743.1	832.6	879.3	814.6	947.7	692.6	634.8	754.2

Table 116. Total mortality (ICD-9 001-799, E800-E999), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	668.9	667.3	670.5	861.8	858.9	864.6	526.3	524.5	528.2
Newfoundland and Labrador	752.4	739.4	765.3	969.9	947.4	992.4	583.6	568.4	598.8
St. John's (NF1)	690.6	669.3	712.0	891.3	853	929.6	540.8	516.1	565.5
Eastern (NF2)	834.3	806.0	862.5	1,130.6	1,080.5	1,180.7	616.2	583.8	648.5
Central (NF3)	752.9	725.2	780.6	934.5	888.7	980.3	601.1	567.3	634.9
Western (NF4)	751.8	719.6	784.0	936.4	882	990.8	602.6	563.6	641.6
Grenfell (NF5)	740.5	659.2	821.9	881.6	751.2	1,011.9	598.5	498.3	698.7
Labrador (NF6)	869.5	757.1	981.8	1,091.3	915.7	1,267.0	652.6	508.2	796.9
Prince Edward Island	695.4	672.4	718.5	941.9	900.6	983.1	513.1	487.2	539.1
PEI/Urban	767.4	732.2	802.7	989.3	926.7	1,051.8	602.6	561.9	643.2
PEI/Rural	633.0	602.8	663.2	902.1	847.4	956.9	433.6	400.5	466.7
Nova Scotia	711.2	702.3	720.1	930.8	914.9	946.8	556.3	545.9	566.7
South-SW (NS1)	711.1	688.8	733.3	919.7	881.2	958.2	550.7	524.6	576.8
Valley (NS2)	641.9	614.6	669.2	834.5	786.6	882.4	507.1	474.8	539.4
Colch-Cumb-E.Hants (NS3)	693.0	668.1	717.9	889.2	845.6	932.8	547.5	518.2	576.8
Pictou-GASHA (NS4)	692.4	666.8	718.0	904.1	859.3	948.9	534.9	505.2	564.7
Cape Breton (NS5)	803.2	779.6	826.8	1,095.7	1,052.1	1,139.3	611.3	584.4	638.1
Capital (NS6)	703.7	688.2	719.2	922.3	893.7	951	558.8	540.9	576.7
New Brunswick	691.4	681.5	701.4	906.4	888.8	924	528.9	517.5	540.4
Moncton (NB1)	637.6	618.6	656.6	848.5	814.6	882.4	477.8	456.2	499.4
Sussex/Saint John (NB2)	722.6	702.1	743.1	936.4	899.6	973.2	575.8	551.9	599.7
Fredericton (NB3)	717.3	695.1	739.4	930.2	891.2	969.2	555.6	529.8	581.3
Edmundston (NB4)	750.1	710.8	789.4	1,003.4	932.9	1,073.9	560.2	515.9	604.5
Campbellton (NB5)	719.6	671.0	768.2	938.9	853.6	1,024.3	547.7	492.2	603.2
Bathurst (NB6)	641.3	612.3	670.2	853.1	803.1	903.2	469.5	436.3	502.7
Miramichi (NB7)	703.9	664.6	743.3	921.9	853.7	990.2	513.9	469.5	558.3

Table 117. Age standardized circulatory disease death rate (ICD-9 390-459), Canada, Atlantic Provinces, and Health Districts, 1996

	All circulatory disease						Ischaemic heart disease						Cerebrovascular disease						All other circulatory disease					
	Both sexes		Male		Female		Both sexes		Male		Female		Both sexes		Male		Female		Both sexes		Male		Female	
	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province
Canada																								
Newfoundland and Labrador	x		x		x		x		x		x		x		x		x		x		x		x	
St. John's (NF1)	x	✓	x	✓	x	✓	x	✓	x	✓	x	✓	x	✓	x	✓	x	✓	x	✓	x	✓	x	✓
Eastern (NF2)	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Central (NF3)	x	x	x	✓	x	x	x	x	x	✓	x	x	x	x	x	x	x	x	✓	x	✓	x	✓	
Western (NF4)	x	✓	x	✓	x	x	x	✓	x	✓	x	x	x	✓	x	✓	x	✓	x	x	x	x	x	x
Grenfell (NF5)	x	x	x	x	x	x	x	x	x	x	x	x	✓	x	✓	✓	✓	x	x	x	x	x	x	x
Labrador (NF6)	x	✓	x	✓	x	✓	✓	✓	✓	✓	✓	✓	x	x	x	x	x	x	x	x	x	x	x	x
Prince Edward Island	x		x		x		x		x		x		x		x		x		x		x		✓	
PEI/Urban	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
PEI/Rural	✓	✓	x	✓	✓	✓	x	✓	x	✓	x	✓	✓	✓	x	✓	✓	✓	✓	✓	x	✓	✓	✓
Nova Scotia	x		x		x		x		x		✓		✓		✓		✓		x		x		x	
South-SW (NS1)	x	x	x	x	x	x	x	x	x	x	x	x	x	✓	x	x	x	x	x	x	x	x	x	x
Valley (NS2)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	x	✓	s	✓	x	x	✓	x	✓	✓	✓	✓
Colch-Cumb-E.Hants (NS3)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	x	✓	x	✓	x	✓
Pictou-GASHA (NS4)	x	✓	x	✓	✓	✓	✓	✓	x	✓	✓	✓	✓	✓	✓	✓	x	x	✓	x	✓	x	✓	
Cape Breton (NS5)	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Capital (NS6)	✓	✓	x	✓	✓	✓	✓	✓	x	✓	✓	✓	✓	✓	✓	✓	✓	✓	x	✓	x	✓	x	x
New Brunswick	x		x		x		✓		✓		x		✓		x		✓		x		x		x	
Moncton (NB1)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	x	x	x	✓	✓	x	x	x	x	x	x	✓
Sussex/Saint John (NB2)	x	x	x	x	x	x	x	x	x	x	x	x	✓	✓	x	x	✓	✓	x	✓	x	✓	x	✓
Fredericton (NB3)	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Edmundston (NB4)	x	x	x	x	x	x	✓	✓	✓	✓	✓	✓	x	x	x	x	✓	x	x	x	x	✓	x	x
Campbellton (NB5)	x	✓	✓	✓	x	✓	✓	✓	✓	✓	✓	x	✓	✓	✓	✓	✓	✓	x	x	x	✓	x	x
Bathurst (NB6)	✓	✓	✓	✓	✓	x	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	x	✓	x	✓	x	x
Miramichi (NB7)	x	x	x	x	x	x	x	x	x	x	x	x	✓	✓	✓	✓	✓	✓	x	✓	x	✓	✓	✓

x - less favourable circulatory disease death rate than the province or nation ✓ - more favourable circulatory disease death rate than the province or nation

s - same circulatory disease death rate as the province or nation

Table 118. All circulatory disease deaths (ICD-9 390-459), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Crude count	95% CI		Crude count	95% CI		Crude count	95% CI	
		low	high		low	high		low	high
Canada	79,283	78,965	79,602	39,945	39,720	40,172	39,338	39,114	39,562
Newfoundland and Labrador	1,719	1,672	1,766	916	883	951	802	771	835
St. John's (NF1)	536	510	563	276	258	296	260	242	278
Eastern (NF2)	457	433	482	240	223	258	217	201	234
Central (NF3)	379	358	402	209	193	226	170	156	185
Western (NF4)	272	254	291	146	132	160	127	114	140
Grenfell (NF5)	48	40	56	28	22	34	20	15	25
Labrador (NF6)	26	21	33	17	13	22	10	7	14
Prince Edward Island	460	436	485	233	217	251	227	210	244
PEI/Urban	238	221	256	110	98	122	128	116	141
PEI/Rural	222	206	240	124	112	137	99	88	110
Nova Scotia	2,938	2,877	2,999	1,487	1,444	1,531	1,451	1,408	1,494
South-SW (NS1)	552	526	579	288	269	307	264	246	283
Valley (NS2)	254	237	273	129	116	142	126	113	139
Colch-Cumb-E.Hants (NS3)	345	324	366	172	158	188	173	158	188
Pictou-GASHA (NS4)	356	335	378	179	165	195	177	162	192
Cape Breton (NS5)	563	536	590	294	275	313	269	251	288
Capital (NS6)	868	835	901	425	402	449	442	419	467
New Brunswick	2,271	2,217	2,325	1,152	1,114	1,190	1,119	1,082	1,157
Moncton (NB1)	543	517	569	271	253	290	272	254	291
Sussex/Saint John (NB2)	609	581	637	299	280	319	309	290	330
Fredericton (NB3)	501	477	527	264	246	282	238	221	256
Edmundston (NB4)	162	148	177	80	71	91	82	72	93
Campbellton (NB5)	95	84	106	45	38	53	50	43	59
Bathurst (NB6)	206	190	223	111	100	124	95	84	106
Miramichi (NB7)	155	141	169	82	72	92	73	64	83

Table 119. All circulatory disease deaths (ICD-9 390-459), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	267.2	266.1	268.3	271.9	270.4	273.4	262.6	261.1	264.1
Newfoundland and Labrador	306.6	298.3	315.0	328.0	315.9	340.4	285.3	274.1	296.9
St. John's (NF1)	287.3	273.6	301.6	301.5	281.5	322.5	273.6	254.9	293.3
Eastern (NF2)	366.9	347.9	386.6	388.1	360.6	417.1	345.9	320.1	373.2
Central (NF3)	334.8	315.8	354.6	366.4	338.6	395.8	302.7	277.3	329.7
Western (NF4)	294.2	274.5	314.8	314.9	286.4	345.4	273.5	247.0	301.9
Grenfell (NF5)	266.3	225.2	312.5	305.8	245.5	376.5	224.8	172.7	287.7
Labrador (NF6)	102.5	81.7	127.0	125.6	94.2	164.1	77.8	53.1	110.1
Prince Edward Island	337.8	320.3	355.9	347.3	322.3	373.7	328.5	304.5	353.9
PEI/Urban	374.2	347.5	402.4	352.9	316.3	392.5	394.6	356.7	435.5
PEI/Rural	305.9	283.4	329.8	342.4	308.9	378.5	269.9	240.4	301.9
Nova Scotia	315.5	308.9	322.1	325.5	316.1	335.1	305.8	296.8	315.0
South-SW (NS1)	432.8	412.3	454.0	456.4	426.8	487.6	409.7	381.9	438.9
Valley (NS2)	304.7	283.6	326.8	312.1	282.1	344.3	297.4	268.6	328.5
Colch-Cumb-E.Hants (NS3)	324.8	305.5	345.0	329.6	302.1	358.9	320.2	293.5	348.6
Pictou-GASHA (NS4)	356.0	335.2	377.8	363.2	333.5	394.8	349.0	320.2	379.6
Cape Breton (NS5)	394.5	376.0	413.6	423.8	396.5	452.4	366.8	342.2	392.7
Capital (NS6)	233.6	224.8	242.7	234.2	221.6	247.3	233.1	220.8	245.9
New Brunswick	301.5	294.5	308.8	308.1	298.0	318.5	295.1	285.2	305.2
Moncton (NB1)	296.8	282.6	311.4	299.0	278.9	320.0	294.6	275.0	315.3
Sussex/Saint John (NB2)	342.2	326.8	358.2	343.4	321.5	366.3	341.1	319.7	363.6
Fredericton (NB3)	303.1	288.1	318.7	319.2	297.6	342.0	287.0	266.6	308.7
Edmundston (NB4)	296.7	271.3	323.9	297.2	261.4	336.4	296.3	261.0	335.0
Campbellton (NB5)	288.3	256.3	323.2	274.7	231.1	324.1	301.6	256.3	352.5
Bathurst (NB6)	230.8	213.1	249.5	247.7	222.2	275.3	213.6	189.8	239.5
Miramichi (NB7)	309.4	282.3	338.5	323.5	284.9	365.8	295.1	257.9	336.1

Table 120. All circulatory disease deaths (ICD-9 390-459), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	245.8	244.8	246.8	316.7	314.9	318.4	192.7	191.6	193.8
Newfoundland and Labrador	320.6	312.0	329.2	406.1	390.9	421.3	252.2	242.3	262.2
St. John's (NF1)	288.9	275.0	302.9	378.0	352.1	403.9	220.6	205.1	236.1
Eastern (NF2)	362.4	343.5	381.2	465.7	432.0	499.3	286.0	264.2	307.7
Central (NF3)	325.5	307.0	344.0	401.1	370.0	432.2	259.7	237.6	281.8
Western (NF4)	318.9	297.5	340.4	386.5	349.9	423.1	261.4	235.7	287.2
Grenfell (NF5)	363.2	304.1	422.3	430.4	337.2	523.6	287.2	215.8	358.6
Labrador (NF6)	289.5	219.7	359.3	366.3	253.1	479.5	228.2	137.5	319.0
Prince Edward Island	267.1	253.0	281.2	356.8	330.8	382.8	198.2	182.8	213.6
PEI/Urban	293.7	272.2	315.3	369.1	329.9	408.3	232.0	208.1	255.8
PEI/Rural	243.7	225.1	262.3	346.2	311.6	380.9	167.9	148.1	187.8
Nova Scotia	259.5	254.1	264.8	345.6	335.6	355.6	197.5	191.5	203.4
South-SW (NS1)	284.1	270.3	297.8	376.2	351.4	401.0	210.7	195.6	225.8
Valley (NS2)	228.9	212.6	245.1	300.1	270.6	329.5	177.2	158.8	195.5
Colch-Cumb-E.Hants (NS3)	237.0	222.5	251.4	307.3	281.2	333.5	186.4	169.9	203.0
Pictou-GASHA (NS4)	251.1	235.9	266.3	328.5	301.0	356.1	189.4	172.7	206.0
Cape Breton (NS5)	306.8	292.4	321.3	438.1	409.4	466.8	224.2	208.5	239.8
Capital (NS6)	245.0	235.7	254.3	322.5	304.8	340.3	191.1	180.8	201.4
New Brunswick	257.7	251.7	263.8	334.3	323.2	345.3	199.1	192.3	205.9
Moncton (NB1)	236.0	224.7	247.4	307.2	286.4	328.1	180.5	168.0	193.1
Sussex/Saint John (NB2)	275.5	263.0	288.1	365.3	341.6	389.0	214.1	200.0	228.3
Fredericton (NB3)	279.0	265.1	293.0	368.1	342.7	393.4	211.3	195.7	226.9
Edmundston (NB4)	270.5	246.9	294.1	338.2	295.8	380.7	219.8	192.6	246.9
Campbellton (NB5)	245.9	217.6	274.2	288.0	239.6	336.3	207.1	174.0	240.2
Bathurst (NB6)	218.1	201.1	235.1	281.5	251.7	311.3	165.9	146.5	185.3
Miramichi (NB7)	269.4	245.3	293.6	346.7	303.8	389.7	204.3	177.3	231.3

Table 121. Ischaemic heart disease deaths (ICD-9 410-414), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Crude count	95% CI		Crude count	95% CI		Crude count	95% CI	
		low	high		low	high		low	high
Canada	43,844	43,608	44,081	24,131	23,955	24,307	19,713	19,555	19,873
Newfoundland and Labrador	950	916	985	547	521	574	403	381	427
St. John's (NF1)	291	273	311	166	152	181	126	113	139
Eastern (NF2)	260	242	279	147	133	161	113	102	126
Central (NF3)	211	195	228	124	112	137	87	77	98
Western (NF4)	149	136	164	86	76	97	63	54	72
Grenfell (NF5)	27	22	34	16	12	21	11	8	16
Labrador (NF6)	11	8	16	8	5	12	3	2	6
Prince Edward Island	250	233	269	134	122	148	116	104	129
PEI/Urban	121	109	134	63	54	72	58	50	67
PEI/Rural	129	117	143	72	63	82	58	50	67
Nova Scotia	1,556	1,512	1,601	894	860	928	662	634	692
South-SW (NS1)	290	271	309	168	154	183	121	109	134
Valley (NS2)	132	120	146	76	67	86	56	48	65
Colch-Cumb-E.Hants (NS3)	175	161	190	96	86	108	79	69	89
Pictou-GASHA (NS4)	185	170	201	108	97	120	77	68	88
Cape Breton (NS5)	318	299	339	186	171	202	133	120	146
Capital (NS6)	455	432	480	259	242	278	196	181	212
New Brunswick	1,129	1,092	1,168	613	586	642	516	491	542
Moncton (NB1)	233	216	251	119	107	132	114	102	127
Sussex/Saint John (NB2)	332	312	353	175	161	191	156	143	171
Fredericton (NB3)	253	236	272	143	130	157	111	99	123
Edmundston (NB4)	75	65	85	42	35	50	33	27	40
Campbellton (NB5)	49	42	58	27	22	33	22	17	28
Bathurst (NB6)	94	84	106	57	49	66	37	31	45
Miramichi (NB7)	93	83	104	50	43	59	43	36	51

Table 122. Ischaemic heart disease deaths (ICD-9 410-414), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	147.8	147.0	148.6	164.2	163.1	165.4	131.6	130.5	132.7
Newfoundland and Labrador	169.5	163.3	175.8	195.7	186.4	205.3	143.4	135.5	151.7
St. John's (NF1)	156.2	146.1	166.8	180.8	165.4	197.2	132.4	119.5	146.3
Eastern (NF2)	208.7	194.5	223.7	236.8	215.5	259.7	180.9	162.5	200.9
Central (NF3)	185.9	171.9	200.8	217.0	195.8	239.9	154.3	136.4	173.9
Western (NF4)	161.3	146.9	176.7	186.6	164.9	210.4	136.0	117.6	156.4
Grenfell (NF5)	152.7	122.2	188.4	174.8	130.3	229.5	129.6	91.1	178.8
Labrador (NF6)	44.1	31.0	60.9	60.3	39.5	88.2	26.8	13.6	47.6
Prince Edward Island	183.8	171.0	197.3	199.9	181.1	220.1	168.1	151.1	186.5
PEI/Urban	190.5	171.7	210.9	201.7	174.3	232.0	179.8	154.6	208.0
PEI/Rural	178.0	160.9	196.3	198.4	173.2	226.3	157.7	135.5	182.5
Nova Scotia	167.1	162.4	171.9	195.6	188.3	203.1	139.6	133.6	145.9
South-SW (NS1)	227.1	212.4	242.6	267.1	244.6	291.1	188.0	169.5	208.1
Valley (NS2)	158.5	143.5	174.7	184.3	161.6	209.4	133.3	114.3	154.6
Colch-Cumb-E.Hants (NS3)	164.8	151.1	179.3	184.3	163.9	206.4	145.9	128.1	165.4
Pictou-GASHA (NS4)	185.3	170.4	201.2	218.8	195.9	243.5	152.8	134.0	173.3
Cape Breton (NS5)	223.2	209.4	237.7	267.9	246.4	290.8	180.9	163.8	199.3
Capital (NS6)	122.6	116.2	129.2	142.8	133.0	153.1	103.3	95.2	111.9
New Brunswick	150.0	145.0	155.1	164.1	156.7	171.7	136.1	129.4	143.0
Moncton (NB1)	127.4	118.2	137.1	131.4	118.3	145.6	123.5	110.9	137.1
Sussex/Saint John (NB2)	186.5	175.2	198.3	201.1	184.5	218.9	172.4	157.3	188.5
Fredericton (NB3)	153.2	142.6	164.3	172.7	156.9	189.7	133.7	119.9	148.6
Edmundston (NB4)	136.5	119.5	155.2	155.4	130.0	184.3	118.0	96.3	143.1
Campbellton (NB5)	149.7	127.0	175.3	166.0	132.7	205.2	133.8	104.5	168.8
Bathurst (NB6)	105.7	93.9	118.5	126.8	108.8	146.9	84.2	69.7	100.9
Miramichi (NB7)	186.1	165.2	208.8	199.4	169.4	233.0	172.5	144.5	204.3

Table 123. Ischaemic heart disease deaths (ICD-9 410-414), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	136.4	135.6	137.1	188.2	186.8	189.5	97.3	96.5	98.1
Newfoundland and Labrador	176.6	170.2	183.1	238.1	226.4	249.8	127.8	120.6	135.0
St. John's (NF1)	157.2	146.8	167.6	225.5	205.4	245.6	107.1	96.2	118.1
Eastern (NF2)	206.6	192.1	221.0	278.1	251.9	304.2	152.2	136	168.4
Central (NF3)	180.1	166.2	194.0	235.8	211.7	259.9	132.9	116.8	149.0
Western (NF4)	171.3	155.5	187.2	217.8	190.6	245.0	131.1	112.5	149.6
Grenfell (NF5)	205.3	160.6	250.0	245.2	174.3	316.0	162.9	108.5	217.4
Labrador (NF6)	116.6	72.1	161.2	165.5	89.1	241.9	74.3	20.3	128.4
Prince Edward Island	147.8	137.1	158.5	203.3	183.5	223.1	102.2	91.0	113.3
PEI/Urban	153.2	137.2	169.1	207.5	177.9	237.1	105.1	88.9	121.3
PEI/Rural	142.8	128.4	157.2	199.2	172.7	225.7	99.1	83.7	114.5
Nova Scotia	139.4	135.4	143.4	204.5	196.8	212.3	92.6	88.5	96.8
South-SW (NS1)	150.9	140.8	161.0	215.7	196.8	234.5	100.7	89.9	111.4
Valley (NS2)	121.8	109.8	133.9	176.4	153.6	199.1	82.9	70.0	95.8
Colch-Cumb-E.Hants (NS3)	121.1	110.7	131.5	169.8	150.3	189.3	86.4	75.0	97.8
Pictou-GASHA (NS4)	134.8	123.4	146.2	197.0	175.5	218.4	86.7	75.0	98.3
Cape Breton (NS5)	175.4	164.3	186.5	270.1	247.5	292.8	111.3	100.1	122.4
Capital (NS6)	129.3	122.5	136.1	191.4	177.8	205.1	86.8	79.7	93.9
New Brunswick	129.6	125.3	134.0	176.0	168.0	184.1	93.9	89.2	98.7
Moncton (NB1)	102.7	95.0	110.3	135.8	121.7	149.9	76.9	68.6	85.3
Sussex/Saint John (NB2)	152.3	142.8	161.8	210.0	192.0	228.1	111.3	100.9	121.7
Fredericton (NB3)	141.8	131.8	151.9	197.6	178.9	216.3	100.1	89.2	111.0
Edmundston (NB4)	124.9	108.6	141.3	173.6	143.1	204.2	89.8	71.9	107.7
Campbellton (NB5)	128.8	108.1	149.5	171.5	134.2	208.9	94.6	71.6	117.7
Bathurst (NB6)	101.7	89.8	113.6	140.9	119.8	162.1	67.1	54.4	79.7
Miramichi (NB7)	161.4	142.6	180.2	212.1	178.3	245.9	117.8	97.3	138.3

Table 124. Cerebrovascular disease deaths (ICD-9 430-438), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Crude count	95% CI		Crude count	95% CI		Crude count	95% CI	
		low	high		low	high		low	high
Canada	15,722	15,580	15,864	6,583	6,492	6,676	9,138	9,031	9,247
Newfoundland and Labrador	335	315	357	154	140	168	182	167	197
St. John's (NF1)	112	100	124	46	39	55	65	57	75
Eastern (NF2)	85	75	96	40	33	48	45	38	53
Central (NF3)	80	71	91	38	32	46	42	35	50
Western (NF4)	44	37	52	21	16	27	23	18	29
Grenfell (NF5)	7	4	10	4	2	7	3	1	5
Labrador (NF6)	7	5	11	4	2	7	3	2	6
Prince Edward Island	94	83	105	41	35	49	52	45	61
PEI/Urban	54	46	63	19	15	25	35	28	42
PEI/Rural	40	33	47	22	17	28	18	13	23
Nova Scotia	526	501	552	200	184	216	326	306	347
South-SW (NS1)	96	86	108	36	29	43	61	52	70
Valley (NS2)	52	44	61	20	16	26	32	26	39
Colch-Cumb-E.Hants (NS3)	65	56	75	26	21	32	39	32	47
Pictou-GASHA (NS4)	64	55	73	22	17	28	42	35	49
Cape Breton (NS5)	101	90	112	38	31	45	63	54	72
Capital (NS6)	148	135	163	58	50	67	90	80	102
New Brunswick	428	405	452	185	170	201	243	226	261
Moncton (NB1)	115	103	127	47	40	55	68	59	77
Sussex/Saint John (NB2)	108	97	120	46	39	55	62	53	71
Fredericton (NB3)	96	85	107	41	34	48	55	47	64
Edmundston (NB4)	32	26	39	15	11	20	16	12	21
Campbellton (NB5)	14	10	18	6	4	9	8	5	11
Bathurst (NB6)	39	32	46	18	14	24	20	16	26
Miramichi (NB7)	26	21	32	11	8	16	15	11	19

Table 125. Cerebrovascular disease (ICD-9 430-438), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	53.0	52.5	53.5	44.8	44.2	45.4	61.0	60.3	61.7
Newfoundland and Labrador	59.8	56.2	63.6	55.0	50.1	60.2	64.6	59.3	70.2
St. John's (NF1)	59.9	53.7	66.5	50.6	42.7	59.5	68.8	59.7	79.0
Eastern (NF2)	68.5	60.5	77.3	64.6	53.8	76.9	72.4	60.9	85.3
Central (NF3)	70.9	62.4	80.3	67.1	55.6	80.2	74.8	62.5	88.7
Western (NF4)	47.5	39.9	56.2	45.4	35.2	57.7	49.7	38.9	62.4
Grenfell (NF5)	37.2	23.4	56.4	43.7	23.7	74.0	30.5	14.2	57.6
Labrador (NF6)	28.5	18.3	42.4	30.1	16.3	51.1	26.8	13.6	47.6
Prince Edward Island	68.8	61.1	77.2	61.5	51.4	73.1	75.8	64.7	88.4
PEI/Urban	85.0	72.7	98.9	62.2	47.7	79.8	106.9	87.8	128.9
PEI/Rural	54.6	45.4	65.1	60.9	47.5	77.0	48.3	36.6	62.7
Nova Scotia	56.5	53.7	59.3	43.8	40.4	47.4	68.7	64.5	73.1
South-SW (NS1)	75.5	67.2	84.6	56.6	46.6	68.1	94.0	81.1	108.4
Valley (NS2)	62.3	53.1	72.6	49.3	38.0	62.9	75.0	61.0	91.2
Colch-Cumb-E.Hants (NS3)	61.2	53.0	70.2	49.7	39.6	61.7	72.3	60.1	86.3
Pictou-GASHA (NS4)	63.7	55.1	73.2	44.6	34.7	56.3	82.3	68.8	97.7
Cape Breton (NS5)	70.6	62.9	78.9	54.8	45.4	65.6	85.5	73.9	98.3
Capital (NS6)	39.9	36.4	43.8	31.9	27.4	36.9	47.6	42.2	53.5
New Brunswick	56.9	53.8	60.1	49.5	45.5	53.7	64.2	59.6	68.9
Moncton (NB1)	62.7	56.3	69.6	51.9	43.9	61.0	73.3	63.7	83.9
Sussex/Saint John (NB2)	60.7	54.4	67.6	53.2	44.8	62.5	68.0	58.7	78.3
Fredericton (NB3)	57.8	51.4	64.8	49.2	41.1	58.6	66.4	56.9	77.1
Edmundston (NB4)	57.9	47.1	70.4	56.7	42.0	74.9	59.0	44.1	77.3
Campbellton (NB5)	41.5	30.1	55.7	36.9	22.5	57.1	45.9	29.8	67.7
Bathurst (NB6)	43.3	36.0	51.7	40.8	31.0	52.7	45.9	35.4	58.5
Miramichi (NB7)	52.0	41.4	64.5	44.9	31.6	62.0	59.3	43.6	78.8

Table 126. Cerebrovascular disease deaths (ICD-9 430-438), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	48.4	47.9	48.8	53.7	52.9	54.5	44.3	43.8	44.8
Newfoundland and Labrador	62.5	58.6	66.4	70.2	63.6	76.7	56.5	51.8	61.3
St. John's (NF1)	59.3	52.9	65.6	63.3	52.6	74.0	54.3	46.6	61.9
Eastern (NF2)	66.8	58.6	75.0	79.4	64.9	93.8	59.0	49.0	69.0
Central (NF3)	69.2	60.5	77.9	75.4	61.5	89.4	63.7	52.6	74.8
Western (NF4)	52.8	43.8	61.8	61.3	45.7	77.0	47.2	36.1	58.3
Grenfell (NF5)	50.6	27.9	73.3	55.1	23.8	86.3	40.5	12.4	68.7
Labrador (NF6)	79.5	44.2	114.7	87.1	33.8	140.4	71.5	26.5	116.6
Prince Edward Island	53.3	47.0	59.7	63.6	52.4	74.8	45.9	38.4	53.4
PEI/Urban	65.3	55.1	75.5	66.0	49.0	83.0	64.4	51.5	77.3
PEI/Rural	43.0	35.2	50.8	61.7	46.8	76.6	29.4	21.1	37.7
Nova Scotia	45.3	43.1	47.5	46.9	43.1	50.7	42.8	40.0	45.5
South-SW (NS1)	48.5	42.8	54.1	47.4	38.4	56.5	45.7	38.9	52.6
Valley (NS2)	45.4	38.2	52.6	46.9	35.2	58.7	43.7	34.6	52.7
Colch-Cumb-E.Hants (NS3)	43.6	37.4	49.8	46.7	36.3	57.1	40.0	32.5	47.5
Pictou-GASHA (NS4)	43.0	36.8	49.2	40.1	30.3	49.8	44.0	36.0	52.0
Cape Breton (NS5)	53.4	47.4	59.4	58.4	47.3	69.4	51.6	44.0	59.1
Capital (NS6)	41.5	37.7	45.4	45.5	38.6	52.4	37.4	32.9	42.0
New Brunswick	47.6	45.0	50.2	54.5	49.9	59.0	42.1	39.0	45.3
Moncton (NB1)	48.5	43.4	53.6	54.0	45.1	63.0	44.2	37.9	50.4
Sussex/Saint John (NB2)	47.4	42.2	52.6	58.9	49.0	68.9	40.6	34.6	46.6
Fredericton (NB3)	52.2	46.2	58.3	57.0	46.8	67.1	47.0	39.7	54.2
Edmundston (NB4)	52.3	41.8	62.7	66.1	46.7	85.5	43.8	31.5	56.1
Campbellton (NB5)	35.0	24.3	45.7	40.3	21.7	58.9	31.5	18.4	44.5
Bathurst (NB6)	40.3	33.0	47.7	45.9	33.7	58.1	35.1	26.1	44.1
Miramichi (NB7)	44.8	34.9	54.8	48.5	32.0	65.0	41.4	28.9	53.8

Table 127. All other circulatory disease deaths (ICD-9 390-409, 415-429, 439-459), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Crude count	95% CI		Crude count	95% CI		Crude count	95% CI	
		low	high		low	high		low	high
Canada	19,717	19,559	19,877	9,231	9,123	9,340	10,486	10,371	10,602
Newfoundland and Labrador	433	410	457	216	200	233	217	201	234
St. John's (NF1)	133	120	147	64	56	74	69	60	79
Eastern (NF2)	112	100	124	54	46	62	58	50	67
Central (NF3)	88	78	99	47	40	55	41	35	49
Western (NF4)	79	69	90	38	32	46	41	34	48
Grenfell (NF5)	14	10	18	8	5	12	6	3	9
Labrador (NF6)	8	5	11	5	3	8	3	1	5
Prince Edward Island	116	104	129	58	50	67	58	50	67
PEI/Urban	63	54	72	28	22	34	35	29	42
PEI/Rural	53	46	62	30	24	37	23	18	29
Nova Scotia	856	823	889	393	371	416	462	438	487
South-SW (NS1)	166	152	181	84	74	94	82	73	93
Valley (NS2)	70	61	80	32	26	39	38	31	45
Colch-Cumb-E.Hants (NS3)	105	94	117	50	42	58	55	47	64
Pictou-GASHA (NS4)	107	96	119	49	42	58	58	50	67
Cape Breton (NS5)	144	131	158	70	61	80	74	64	84
Capital (NS6)	264	246	283	108	97	120	156	142	171
New Brunswick	713	683	744	353	333	375	360	339	382
Moncton (NB1)	195	180	211	105	94	117	90	80	102
Sussex/Saint John (NB2)	169	155	184	78	68	88	91	81	103
Fredericton (NB3)	152	139	167	80	71	91	72	63	82
Edmundston (NB4)	56	48	65	23	18	29	33	27	40
Campbellton (NB5)	32	26	39	12	8	16	20	16	26
Bathurst (NB6)	73	64	83	36	30	43	37	31	44
Miramichi (NB7)	36	29	43	20	15	26	16	12	21

Table 128. All other circulatory disease deaths (ICD-9 390-409, 415-429, 439-459), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	66.5	65.9	67.0	62.8	62.1	63.6	70.0	69.2	70.8
Newfoundland and Labrador	77.3	73.2	81.6	77.3	71.5	83.4	77.3	71.5	83.4
St. John's (NF1)	71.3	64.6	78.5	70.2	60.8	80.6	72.4	63.0	82.7
Eastern (NF2)	89.6	80.4	99.6	86.7	74.0	100.8	92.6	79.6	107.1
Central (NF3)	78.0	69.0	87.8	82.3	69.5	96.7	73.6	61.5	87.4
Western (NF4)	85.3	75.0	96.7	82.9	68.7	99.1	87.8	73.2	104.4
Grenfell (NF5)	76.3	55.5	102.5	87.4	57.3	127.8	64.8	39.0	101.4
Labrador (NF6)	29.8	19.4	44.0	35.2	20.0	57.4	24.1	11.8	44.1
Prince Edward Island	85.2	76.6	94.5	85.8	73.7	99.3	84.5	72.7	97.8
PEI/Urban	98.7	85.3	113.5	89.0	71.4	109.8	107.9	88.7	130.0
PEI/Rural	73.4	62.7	85.4	83.1	67.2	101.6	63.8	50.1	80.1
Nova Scotia	91.9	88.4	95.5	86.1	81.3	91.1	97.5	92.4	102.7
South-SW (NS1)	130.1	119.1	141.9	132.8	117.1	149.9	127.6	112.4	144.3
Valley (NS2)	83.9	73.1	95.8	78.4	63.9	95.2	89.2	73.8	106.7
Colch-Cumb-E.Hants (NS3)	98.9	88.4	110.2	95.6	81.2	111.9	102.0	87.3	118.4
Pictou-GASHA (NS4)	107.0	95.8	119.2	99.9	84.8	117.0	113.9	97.9	131.8
Cape Breton (NS5)	100.7	91.6	110.6	101.0	88.0	115.4	100.5	87.8	114.4
Capital (NS6)	71.1	66.3	76.2	59.5	53.2	66.2	82.2	75.0	89.9
New Brunswick	94.7	90.7	98.8	94.5	89.0	100.3	94.8	89.3	100.6
Moncton (NB1)	106.6	98.3	115.5	115.6	103.3	128.9	97.9	86.7	110.0
Sussex/Saint John (NB2)	95.0	87.0	103.6	89.1	78.2	101.1	100.7	89.3	113.2
Fredericton (NB3)	92.1	83.9	100.8	97.3	85.5	110.1	87.0	75.9	99.1
Edmundston (NB4)	102.4	87.7	118.7	85.1	66.7	107.0	119.2	97.4	144.5
Campbellton (NB5)	97.1	79.1	118.0	71.7	50.7	98.6	121.8	94.0	155.3
Bathurst (NB6)	81.8	71.5	93.1	80.1	66.0	96.3	83.5	69.0	100.1
Miramichi (NB7)	71.4	58.8	85.8	79.2	61.0	101.2	63.3	47.1	83.4

Table 129. All other circulatory disease deaths (ICD-9 390-409, 415-429, 439-459), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	61.0	60.5	61.5	74.8	73.9	75.7	51.1	50.6	51.7
Newfoundland and Labrador	81.5	77.0	85.9	97.8	90.1	105.5	67.9	62.7	73.1
St. John's (NF1)	72.5	65.4	79.6	89.2	76.2	102.2	59.2	51.0	67.4
Eastern (NF2)	89.0	79.5	98.5	108.3	91.2	125.3	74.7	63.6	85.9
Central (NF3)	76.2	67.1	85.4	89.9	74.8	104.9	63.0	52.0	74.1
Western (NF4)	94.7	82.7	106.7	107.4	87.1	127.6	83.2	68.5	97.8
Grenfell (NF5)	107.3	74.0	140.7	130.2	76.1	184.3	83.7	44.0	123.4
Labrador (NF6)	93.4	51.7	135.2	113.7	47.2	180.3	82.4	23.4	141.4
Prince Edward Island	65.9	58.9	73.0	89.9	76.5	103.3	50.2	42.4	57.9
PEI/Urban	75.3	64.4	86.2	95.6	75.1	116.1	62.4	50.0	74.8
PEI/Rural	57.9	48.8	67.0	85.4	67.7	103.0	39.4	29.7	49.0
Nova Scotia	74.8	71.9	77.7	94.2	88.8	99.6	62.1	58.7	65.4
South-SW (NS1)	84.7	77.1	92.2	113.1	99.1	127.2	64.3	55.9	72.7
Valley (NS2)	61.6	53.2	70.0	76.8	61.5	92.0	50.6	40.9	60.3
Colch-Cumb-E.Hants (NS3)	72.3	64.2	80.3	90.8	76.3	105.3	60.0	50.4	69.6
Pictou-GASHA (NS4)	73.3	65.2	81.5	91.5	76.7	106.3	58.7	49.6	67.8
Cape Breton (NS5)	78.0	70.6	85.4	109.6	94.4	124.8	61.3	53.0	69.6
Capital (NS6)	74.2	69.0	79.3	85.6	76.1	95.1	66.9	60.7	73.0
New Brunswick	80.5	77.1	84.0	103.8	97.5	110.0	63.1	59.2	66.9
Moncton (NB1)	84.9	78.0	91.8	117.4	104.4	130.4	59.4	52.2	66.7
Sussex/Saint John (NB2)	75.9	69.2	82.5	96.3	83.8	108.9	62.3	54.6	70.0
Fredericton (NB3)	85.0	77.2	92.8	113.5	99.1	127.9	64.2	55.5	73.0
Edmundston (NB4)	93.3	79.3	107.3	98.5	75.0	122.0	86.2	69.3	103.0
Campbellton (NB5)	82.1	65.7	98.5	76.1	50.8	101.4	81.0	60.6	101.4
Bathurst (NB6)	76.1	66.0	86.1	94.6	76.8	112.3	63.8	51.7	75.8
Miramichi (NB7)	63.2	51.2	75.1	86.1	64.2	108.0	45.2	32.0	58.3

Table 130. Age standardized unintentional injury death rate (ICD-9 E800-E929, excluding 870-879), Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes		Male		Female	
	Canada	Province	Canada	Province	Canada	Province
Canada						
Newfoundland and Labrador	✓		✓		✓	
St. John's (NF1)	✓	✓	✓	✓	✓	✓
Eastern (NF2)	✓	×	×	×	✓	×
Central (NF3)	✓	×	✓	×	✓	×
Western (NF4)	✓	×	×	×	✓	✓
Grenfell (NF5)	×	×	✓	×	×	×
Labrador (NF6)	×	×	×	×	×	×
Prince Edward Island	×		×		✓	
PEI/Urban	×	✓	×	✓	✓	✓
PEI/Rural	×	×	×	×	×	×
Nova Scotia	×		×		✓	
South-SW (NS1)	×	×	×	×	×	×
Valley (NS2)	✓	✓	✓	✓	✓	✓
Colch-Cumb-E.Hants (NS3)	×	×	×	×	×	×
Pictou-GASHA (NS4)	×	×	×	×	✓	×
Cape Breton (NS5)	×	×	×	×	✓	✓
Capital (NS6)	✓	✓	✓	✓	✓	✓
New Brunswick	×		×		✓	
Moncton (NB1)	✓	✓	×	✓	✓	✓
Sussex/Saint John (NB2)	✓	✓	✓	✓	×	×
Fredericton (NB3)	×	✓	×	×	✓	✓
Edmundston (NB4)	×	×	×	×	×	×
Campbellton (NB5)	×	×	×	×	×	×
Bathurst (NB6)	✓	✓	×	✓	✓	✓
Miramichi (NB7)	×	×	×	×	×	×

× - less favourable unintentional injury death rate than the province or nation ✓ - more favourable unintentional injury death rate than the province or nation

s - same unintentional injury death rate as the province or nation

Table 131. Unintentional injury deaths (ICD-9 E800-E929, excluding 870-879), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Crude count	95% CI		Crude count	95% CI		Crude count	95% CI	
		low	high		low	high		low	high
Canada	8,460	8,356	8,564	5,285	5,203	5,368	3,175	3,111	3,239
Newfoundland and Labrador	131	119	144	92	82	104	39	32	46
St. John's (NF1)	30	24	37	18	13	23	12	9	17
Eastern (NF2)	35	28	42	25	20	32	9	6	13
Central (NF3)	30	24	36	21	17	27	8	6	12
Western (NF4)	21	17	27	18	13	23	4	2	6
Grenfell (NF5)	5	3	8	3	1	5	2	1	4
Labrador (NF6)	10	7	14	7	5	11	3	1	5
Prince Edward Island	50	42	58	34	28	41	16	12	21
PEI/Urban	20	16	26	14	11	19	6	4	9
PEI/Rural	29	24	36	19	15	25	10	7	14
Nova Scotia	290	272	310	183	168	199	107	96	119
South-SW (NS1)	50	42	58	32	26	39	18	13	23
Valley (NS2)	24	19	30	14	11	19	10	7	14
Colch-Cumb-E.Hants (NS3)	42	35	49	25	20	31	17	13	22
Pictou-GASHA (NS4)	37	30	44	23	18	29	13	10	18
Cape Breton (NS5)	47	40	55	29	24	36	18	13	23
Capital (NS6)	91	81	103	60	51	69	32	26	39
New Brunswick	240	223	258	160	146	175	80	70	91
Moncton (NB1)	55	47	64	35	29	43	20	15	25
Sussex/Saint John (NB2)	51	43	59	29	23	36	22	17	27
Fredericton (NB3)	51	43	59	36	29	43	15	11	20
Edmundston (NB4)	26	21	32	20	15	26	6	4	9
Campbellton (NB5)	13	10	18	9	6	13	4	2	7
Bathurst (NB6)	25	20	32	19	14	24	7	4	10
Miramichi (NB7)	19	15	25	13	9	17	7	4	10

Table 132. Unintentional injury deaths (ICD-9 E800-E929, excluding 870-879), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	28.5	28.2	28.9	36.0	35.4	36.5	21.2	20.8	21.6
Newfoundland and Labrador	23.4	21.1	25.8	33.0	29.3	37.1	13.8	11.4	16.4
St. John's (NF1)	16.1	13.0	19.7	19.3	14.6	25.0	13.0	9.3	17.7
Eastern (NF2)	27.8	22.8	33.6	40.9	32.4	50.9	14.9	10.1	21.2
Central (NF3)	26.2	21.1	32.1	37.3	29.0	47.3	14.8	9.8	21.5
Western (NF4)	23.0	17.9	29.2	38.2	28.9	49.5	7.9	4.2	13.7
Grenfell (NF5)	27.9	16.2	44.9	32.8	16.0	59.9	22.9	9.3	47.3
Labrador (NF6)	40.2	27.8	56.3	55.3	35.5	82.2	24.1	11.8	44.1
Prince Edward Island	36.5	31.0	42.7	50.1	41.0	60.6	23.2	17.3	30.5
PEI/Urban	32.0	24.7	40.8	46.1	33.8	61.5	18.5	11.3	28.6
PEI/Rural	40.4	32.6	49.5	53.5	41.0	68.7	27.4	18.8	38.5
Nova Scotia	31.2	29.2	33.3	40.1	36.9	43.6	22.6	20.2	25.1
South-SW (NS1)	38.9	33.0	45.6	50.8	41.3	61.7	27.4	20.7	35.5
Valley (NS2)	28.7	22.7	36.0	34.8	25.5	46.4	22.9	15.6	32.4
Colch-Cumb-E.Hants (NS3)	39.2	32.8	46.6	47.2	37.3	58.9	31.5	23.7	41.1
Pictou-GASHA (NS4)	36.7	30.3	44.0	47.3	37.1	59.3	26.3	19.1	35.5
Cape Breton (NS5)	33.0	27.8	38.7	42.3	34.1	51.9	24.1	18.2	31.3
Capital (NS6)	24.6	21.8	27.6	32.8	28.3	37.9	16.7	13.6	20.3
New Brunswick	31.9	29.6	34.3	42.9	39.2	46.9	21.1	18.6	23.9
Moncton (NB1)	30.1	25.7	34.9	39.0	32.1	47.0	21.3	16.4	27.3
Sussex/Saint John (NB2)	28.5	24.2	33.3	33.3	26.8	40.8	23.9	18.6	30.2
Fredericton (NB3)	30.6	26.0	35.8	43.2	35.6	52.0	18.1	13.4	24.0
Edmundston (NB4)	47.5	37.8	59.0	74.0	56.9	94.5	21.7	13.2	33.5
Campbellton (NB5)	40.5	29.3	54.5	55.3	37.2	79.3	26.0	14.4	43.2
Bathurst (NB6)	28.4	22.5	35.3	41.5	31.7	53.5	15.0	9.4	22.8
Miramichi (NB7)	38.7	29.6	49.6	50.2	36.0	68.1	27.0	16.9	40.8

Table 133. Unintentional injury deaths (ICD-9 E800-E929, excluding 870-879), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	27.7	27.3	28.0	38.3	37.7	38.9	17.6	17.2	18.0
Newfoundland and Labrador	23.6	21.2	25.9	34.1	29.9	38.2	12.6	10.3	14.9
St. John's (NF1)	16.0	12.6	19.3	21.4	15.4	27.5	10.9	7.3	14.4
Eastern (NF2)	27.0	21.8	32.3	40.5	31.2	49.8	13.1	8.2	18.0
Central (NF3)	26.0	20.6	31.5	37.3	28.1	46.6	13.0	7.9	18.2
Western (NF4)	24.0	18.0	30.0	39.8	28.7	50.9	8.0	3.3	12.8
Grenfell (NF5)	30.2	14.6	45.7	34.3	11.6	56.9	24.8	4.7	44.9
Labrador (NF6)	52.5	29.6	75.3	71.0	31.7	110.3	34.7	8.7	60.7
Prince Edward Island	33.7	28.2	39.2	51.4	41.3	61.6	17.4	12.2	22.6
PEI/Urban	29.6	22.0	37.2	47.4	33.1	61.7	13.2	6.8	19.7
PEI/Rural	37.5	29.5	45.5	55.0	40.8	69.3	21.1	13.1	29.1
Nova Scotia	28.6	26.7	30.6	41.5	38.0	45.0	16.9	15.0	18.9
South-SW (NS1)	35.1	29.2	40.9	50.3	40.1	60.6	20.6	14.5	26.6
Valley (NS2)	24.1	18.4	29.7	33.5	23.4	43.5	15.1	9.3	20.9
Colch-Cumb-E.Hants (NS3)	33.6	27.5	39.7	47.8	36.8	58.9	22.9	16.2	29.6
Pictou-GASHA (NS4)	30.9	24.9	36.9	45.7	34.8	56.7	17.2	11.3	23.1
Cape Breton (NS5)	30.3	25.2	35.4	43.7	34.4	53.1	16.7	11.9	21.5
Capital (NS6)	24.9	21.9	27.9	36.1	30.7	41.5	14.4	11.5	17.4
New Brunswick	29.9	27.7	32.1	43.8	39.9	47.8	16.8	14.6	19.1
Moncton (NB1)	27.1	22.9	31.3	39.6	32.0	47.2	16.6	12.2	21.1
Sussex/Saint John (NB2)	25.6	21.4	29.7	35.1	27.6	42.7	17.7	13.2	22.3
Fredericton (NB3)	29.6	24.8	34.3	44.2	35.7	52.6	15.5	10.9	20.2
Edmundston (NB4)	46.9	36.4	57.4	78.7	58.4	99.0	18.5	9.8	27.2
Campbellton (NB5)	37.3	25.6	48.9	54.3	33.7	75.0	21.2	9.3	33.2
Bathurst (NB6)	27.2	21.0	33.4	41.0	30.1	51.9	13.2	7.2	19.2
Miramichi (NB7)	36.6	27.1	46.2	49.6	33.7	65.4	20.6	11.1	30.0

Table 134. Age standardized AIDS death rate (ICD-9 042-044), Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes		Male		Female	
	Canada	Province	Canada	Province	Canada	Province
Canada						
Newfoundland and Labrador	✓		✓		-	
St. John's (NF1)	✓	×	✓	×	-	-
Eastern (NF2)	✓	✓	-	-	-	-
Central (NF3)	-	-	-	-	-	-
Western (NF4)	✓	×	✓	×	-	-
Grenfell (NF5)	-	-	-	-	-	-
Labrador (NF6)	-	-	-	-	-	-
Prince Edward Island	✓		✓		-	
PEI/Urban	-	-	-	-	-	-
PEI/Rural	-	-	-	-	-	-
Nova Scotia	✓		✓			
South-SW (NS1)	-	-	-	-	-	-
Valley (NS2)	-	-	-	-	-	-
Colch-Cumb-E.Hants (NS3)	-	-	-	-	-	-
Pictou-GASHA (NS4)	-	-	-	-	-	-
Cape Breton (NS5)	✓	✓	✓	✓	-	-
Capital (NS6)	✓	×	✓	×		-
New Brunswick	✓		✓		✓	
Moncton (NB1)	✓	×	✓	×	-	-
Sussex/Saint John (NB2)	✓	s	✓	×	-	-
Fredericton (NB3)	✓	✓	✓	✓	-	-
Edmundston (NB4)	-	-	-	-	-	-
Campbellton (NB5)	-	-	-	-	-	-
Bathurst (NB6)	-	-	-	-	-	-
Miramichi (NB7)	-	-	-	-	-	-

× - less favourable AIDS death rate than the province or nation

✓ - more favourable AIDS death rate than the province or nation

s - same AIDS death rate as the province or nation

“-” - no data available for comparison

Table 135. AIDS deaths (ICD-9 042-044), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Crude count	95% CI		Crude count	95% CI		Crude count	95% CI	
		low	high		low	high		low	high
Canada	1,232	1,192	1,272	1,129	1,091	1,167	103	92	115
Newfoundland and Labrador	7	5	11	7	4	10	1	--	2
St. John's (NF1)	4	2	6	4	2	6	--	--	--
Eastern (NF2)	1	--	3	1	--	2	1	--	2
Central (NF3)	1	--	3	1	--	3	--	--	--
Western (NF4)	1	--	3	1	--	3	--	--	--
Grenfell (NF5)	--	--	--	--	--	--	--	--	--
Labrador (NF6)	--	--	--	--	--	--	--	--	--
Prince Edward Island	1	--	3	1	--	3	--	--	--
PEI/Urban	1	--	2	1	--	2	--	--	--
PEI/Rural	1	--	2	1	--	2	--	--	--
Nova Scotia	19	14	24	18	14	23	1	--	2
South-SW (NS1)	1	--	3	1	--	3	--	--	--
Valley (NS2)	1	--	3	1	--	3	--	--	--
Colch-Cumb-E.Hants (NS3)	1	--	3	1	--	2	--	--	2
Pictou-GASHA (NS4)	1	--	3	1	--	3	--	--	2
Cape Breton (NS5)	1	--	3	1	--	3	--	--	--
Capital (NS6)	13	9	18	13	9	18	--	--	--
New Brunswick	14	10	19	13	9	17	2	1	4
Moncton (NB1)	7	4	10	6	4	10	--	--	2
Sussex/Saint John (NB2)	3	2	6	3	2	6	--	--	--
Fredericton (NB3)	2	1	4	1	--	3	--	--	2
Edmundston (NB4)	1	--	3	--	--	2	1	--	2
Campbellton (NB5)	--	--	2	--	--	2	--	--	--
Bathurst (NB6)	1	--	2	1	--	2	--	--	--
Miramichi (NB7)	1	--	2	--	--	2	--	--	2

Table 136. AIDS deaths (ICD-9 042-044), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	4.2	4.0	4.3	7.7	7.4	7.9	0.7	0.6	0.8
Newfoundland and Labrador	1.3	0.8	1.9	2.4	1.5	3.6	0.2	--	0.8
St. John's (NF1)	2.0	1.0	3.4	4.0	2.1	6.9	--	--	--
Eastern (NF2)	1.1	0.3	2.6	1.1	0.2	3.5	1.1	0.2	3.5
Central (NF3)	0.9	0.2	2.4	1.8	0.5	4.7	--	--	--
Western (NF4)	1.4	0.5	3.4	2.9	0.9	6.9	--	--	--
Grenfell (NF5)	--	--	--	--	--	--	--	--	--
Labrador (NF6)	--	--	--	--	--	--	--	--	--
Prince Edward Island	1.0	0.3	2.3	2.0	0.6	4.7	--	--	--
PEI/Urban	1.0	0.2	3.4	2.1	0.4	7.0	--	--	--
PEI/Rural	0.9	0.2	3.0	1.8	0.3	6.0	--	--	--
Nova Scotia	2.0	1.5	2.6	3.9	3.0	5.1	0.1	--	0.5
South-SW (NS1)	0.8	0.2	2.1	1.6	0.4	4.3	--	--	--
Valley (NS2)	1.2	0.3	3.2	2.4	0.6	6.5	--	--	--
Colch-Cumb-E.Hants (NS3)	0.9	0.2	2.5	1.3	0.2	4.1	0.6	--	3.0
Pictou-GASHA (NS4)	1.3	0.4	3.2	2.0	0.5	5.4	0.7	--	3.2
Cape Breton (NS5)	0.9	0.3	2.2	1.9	0.6	4.6	--	--	--
Capital (NS6)	3.5	2.5	4.7	7.2	5.2	9.7	--	--	--
New Brunswick	1.9	1.4	2.5	3.4	2.4	4.6	0.4	0.2	1.0
Moncton (NB1)	3.6	2.3	5.5	7.0	4.3	10.7	0.4	--	1.7
Sussex/Saint John (NB2)	1.9	1.0	3.3	3.8	1.9	6.8	--	--	--
Fredericton (NB3)	1.0	0.4	2.2	1.6	0.5	3.9	0.4	--	1.9
Edmundston (NB4)	1.8	0.5	4.9	1.2	0.1	5.9	2.4	0.4	7.8
Campbellton (NB5)	1.0	0.1	4.8	2.0	0.1	9.8	--	--	--
Bathurst (NB6)	0.7	0.1	2.4	1.5	0.3	4.8	--	--	--
Miramichi (NB7)	1.3	0.2	4.3	1.3	0.1	6.3	1.3	0.1	6.5

Table 137. AIDS deaths (ICD-9 042-044), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	4.0	3.9	4.2	7.4	7.1	7.6	0.7	0.6	0.8
Newfoundland and Labrador	1.4	0.8	2.0	2.5	1.4	3.6	--	--	--
St. John's (NF1)	2.0	0.8	3.1	4.0	1.6	6.3	--	--	--
Eastern (NF2)	1.2	--	2.4	--	--	--	--	--	--
Central (NF3)	--	--	--	--	--	--	--	--	--
Western (NF4)	1.6	--	3.2	3.3	--	6.5	--	--	--
Grenfell (NF5)	--	--	--	--	--	--	--	--	--
Labrador (NF6)	--	--	--	--	--	--	--	--	--
Prince Edward Island	1.0	--	2.0	2.0	--	4.0	--	--	--
PEI/Urban	--	--	--	--	--	--	--	--	--
PEI/Rural	--	--	--	--	--	--	--	--	--
Nova Scotia	2.0	1.5	2.5	3.8	2.8	4.8	--	--	--
South-SW (NS1)	--	--	--	--	--	--	--	--	--
Valley (NS2)	--	--	--	--	--	--	--	--	--
Colch-Cumb-E.Hants (NS3)	--	--	--	--	--	--	--	--	--
Pictou-GASHA (NS4)	--	--	--	--	--	--	--	--	--
Cape Breton (NS5)	1.1	--	2.1	2.2	--	4.4	--	--	--
Capital (NS6)	3.2	2.2	4.2	6.6	4.5	8.7	--	--	--
New Brunswick	1.9	1.3	2.5	3.3	2.3	4.4	0.5	--	0.9
Moncton (NB1)	3.5	2.0	5.1	6.7	3.7	9.8	--	--	--
Sussex/Saint John (NB2)	1.9	0.7	3.0	3.7	1.4	6.1	--	--	--
Fredericton (NB3)	1.1	0.1	2.0	1.6	--	3.3	--	--	--
Edmundston (NB4)	--	--	--	--	--	--	--	--	--
Campbellton (NB5)	--	--	--	--	--	--	--	--	--
Bathurst (NB6)	--	--	--	--	--	--	--	--	--
Miramichi (NB7)	--	--	--	--	--	--	--	--	--

Table 138. Age standardized cancer (malignant neoplasms) death rate (ICD-9 140-208), Canada, Atlantic Provinces, and Health Districts, 1996

	All cancer						Lung cancer						Colorectal cancer						Prostate cancer		Female breast cancer	
	Both sexes		Male		Female		Both sexes		Male		Female		Both sexes		Male		Female		Canada	Province	Canada	Province
	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province
Canada																						
Newfoundland and Labrador	✓		×		✓		✓		×		✓		s		✓		×		×		✓	
St. John's (NF1)	✓	×	s	✓	×	×	✓	✓	×	✓	×	✓	×	✓	✓	✓	s	✓	✓	✓	×	×
Eastern (NF2)	✓	✓	×	×	✓	✓	✓	✓	×	×	✓	✓	✓	×	×	×	×	×	×	×	✓	✓
Central (NF3)	×	×	✓	✓	✓	✓	×	×	✓	✓	✓	✓	×	✓	✓	✓	×	×	✓	✓	×	×
Western (NF4)	✓	×	×	✓	✓	✓	×	×	×	✓	✓	✓	s	s	✓	s	×	✓	×	×	×	×
Grenfell (NF5)	×	×	✓	✓	✓	✓	×	×	×	×	-	-	×	×	✓	✓	s	✓	-	-	×	×
Labrador (NF6)	✓	✓	-	-	-	-	✓	✓	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Prince Edward Island	✓		×		✓		✓		×		×		×		✓		×		×		✓	
PEI/Urban	✓	✓	×	✓	×	×	✓	×	✓	✓	×	×	×	×	✓	×	×	×	×	×	✓	×
PEI/Rural	×	×	×	×	✓	✓	✓	✓	×	×	×	✓	×	s	✓	✓	✓	✓	×	×	✓	✓
Nova Scotia	✓		×		×		✓		×		×		×		✓		✓		×		×	
South-SW (NS1)	✓	×	×	✓	×	✓	✓	×	✓	✓	×	✓	×	×	✓	✓	×	×	s	✓	×	✓
Valley (NS2)	✓	×	×	✓	×	✓	✓	×	×	✓	×	✓	✓	✓	×	×	×	×	×	×	✓	✓
Colch-Cumb-E.Hants (NS3)	✓	×	×	✓	×	✓	✓	✓	×	×	×	✓	×	×	✓	✓	✓	s	×	✓	✓	✓
Pictou-GASHA (NS4)	✓	✓	×	×	×	×	✓	s	×	×	×	✓	×	×	✓	✓	✓	✓	✓	✓	×	✓
Cape Breton (NS5)	✓	✓	×	×	×	×	✓	✓	×	×	×	×	×	✓	×	×	✓	✓	×	×	×	×
Capital (NS6)	✓	×	×	✓	×	×	✓	×	×	✓	×	×	×	×	✓	✓	✓	✓	×	×	×	×
New Brunswick	✓		×		×		✓		×		✓		×		✓		✓		×		×	
Moncton (NB1)	✓	✓	×	×	×	s	✓	✓	×	✓	×	×	×	✓	✓	×	✓	×	×	×	×	×
Sussex/Saint John (NB2)	✓	✓	×	✓	×	×	✓	✓	×	✓	×	×	×	×	✓	✓	✓	×	✓	✓	×	×
Fredericton (NB3)	×	×	×	✓	✓	✓	×	×	×	✓	✓	✓	×	✓	✓	✓	✓	×	✓	✓	✓	✓
Edmundston (NB4)	✓	✓	×	×	✓	✓	✓	✓	×	×	✓	✓	✓	✓	✓	×	×	×	✓	✓	✓	✓
Campbellton (NB5)	✓	✓	×	×	✓	✓	✓	✓	×	×	×	×	×	×	-	-	-	-	✓	✓	×	×
Bathurst (NB6)	✓	×	×	✓	✓	✓	✓	×	×	✓	✓	✓	×	✓	×	×	✓	✓	×	×	×	×
Miramichi (NB7)	✓	✓	×	×	×	×	✓	✓	×	✓	×	×	×	×	✓	✓	✓	✓	✓	✓	✓	✓

× - less favourable cancer death rate than the province or nation ✓ - more favourable cancer death rate than the province or nation
 s - same cancer death rate as the province or nation “-” - no data available for comparison

Table 139. All cancer deaths (malignant neoplasms) (ICD-9 140-208), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Crude count	95% CI		Crude count	95% CI		Crude count	95% CI	
		low	high		low	high		low	high
Canada	58,570	58,297	58,844	31,507	31,307	31,708	27,063	26,877	27,249
Newfoundland and Labrador	1,057	1,021	1,094	608	581	637	449	425	473
St. John's (NF1)	344	323	365	184	169	200	160	146	174
Eastern (NF2)	278	260	298	175	161	190	103	92	115
Central (NF3)	215	199	232	122	110	135	94	83	105
Western (NF4)	170	156	186	98	87	110	72	63	82
Grenfell (NF5)	24	19	30	13	9	18	11	8	16
Labrador (NF6)	25	20	31	16	12	21	8	6	12
Prince Edward Island	300	281	320	172	158	187	128	116	142
PEI/Urban	147	134	161	78	68	88	69	60	79
PEI/Rural	153	140	168	94	84	106	59	51	68
Nova Scotia	2,215	2,162	2,269	1,174	1,136	1,213	1,041	1,005	1,078
South-SW (NS1)	348	328	370	183	168	198	166	152	181
Valley (NS2)	198	183	214	109	97	121	89	79	100
Colch-Cumb-E.Hants (NS3)	273	255	292	146	133	160	127	114	140
Pictou-GASHA (NS4)	269	251	288	147	134	162	122	110	135
Cape Breton (NS5)	406	383	429	221	205	238	185	170	201
Capital (NS6)	721	691	752	368	347	390	353	332	375
New Brunswick	1,633	1,588	1,679	914	881	949	719	689	750
Moncton (NB1)	417	394	441	235	218	252	182	168	198
Sussex/Saint John (NB2)	409	387	432	214	198	231	195	180	212
Fredericton (NB3)	318	299	339	175	160	190	144	131	158
Edmundston (NB4)	118	106	131	73	63	83	46	38	54
Campbellton (NB5)	80	70	90	49	41	57	31	25	38
Bathurst (NB6)	174	160	190	100	89	112	74	65	84
Miramichi (NB7)	117	105	129	70	61	80	47	40	55

Table 140. All cancer deaths (malignant neoplasms) (ICD-9 140-208), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	197.4	196.5	198.3	214.5	213.1	215.8	180.7	179.4	181.9
Newfoundland and Labrador	188.5	182.1	195.2	217.7	207.9	227.9	159.5	151.2	168.2
St. John's (NF1)	184.4	173.4	195.9	201.1	184.9	218.4	168.2	153.7	183.8
Eastern (NF2)	223.4	208.7	239.0	282.6	259.2	307.5	165.0	147.4	184.1
Central (NF3)	190.1	175.8	205.1	213.0	191.9	235.6	166.8	148.1	187.1
Western (NF4)	184.0	168.5	200.4	211.8	188.7	237.1	156.2	136.4	178.0
Grenfell (NF5)	135.9	107.3	169.8	142.0	102.4	191.9	129.6	91.1	178.8
Labrador (NF6)	96.0	75.9	119.8	123.1	92.0	161.2	67.1	44.3	97.4
Prince Edward Island	220.5	206.5	235.3	256.0	234.6	278.8	186.0	168.1	205.3
PEI/Urban	231.5	210.6	253.8	249.9	219.4	283.5	213.8	186.2	244.3
PEI/Rural	211.0	192.4	230.9	261.2	232.1	292.9	161.4	138.9	186.4
Nova Scotia	237.9	232.2	243.6	257.0	248.6	265.6	219.4	211.8	227.2
South-SW (NS1)	273.1	256.9	290.0	289.8	266.3	314.8	256.8	234.9	280.0
Valley (NS2)	237.2	218.7	256.8	263.6	236.1	293.3	211.4	187.2	237.9
Colch-Cumb-E.Hants (NS3)	257.0	239.9	275.1	279.9	254.6	306.9	234.9	212.2	259.4
Pictou-GASHA (NS4)	269.0	250.9	288.0	298.4	271.6	327.2	240.3	216.6	265.9
Cape Breton (NS5)	284.4	268.8	300.7	318.9	295.4	343.9	251.8	231.5	273.4
Capital (NS6)	194.1	186.1	202.4	202.6	190.9	214.8	186.0	175.1	197.5
New Brunswick	216.9	210.9	223.0	244.6	235.6	253.9	189.6	181.7	197.7
Moncton (NB1)	228.1	215.7	240.9	259.2	240.6	278.8	197.5	181.5	214.6
Sussex/Saint John (NB2)	230.0	217.4	243.1	245.1	226.7	264.6	215.4	198.5	233.4
Fredericton (NB3)	192.5	180.6	205.0	211.5	193.9	230.1	173.5	157.7	190.5
Edmundston (NB4)	216.3	194.7	239.6	268.8	234.9	306.2	165.0	139.1	194.4
Campbellton (NB5)	241.8	212.6	273.9	299.3	253.6	350.7	185.8	150.8	226.4
Bathurst (NB6)	195.3	179.1	212.6	223.2	199.1	249.5	167.0	146.1	190.0
Miramichi (NB7)	233.4	209.9	258.8	275.9	240.4	315.2	190.0	160.5	223.3

Table 141. All cancer deaths (malignant neoplasms) (ICD-9 140-208), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	185.7	184.8	186.5	235.3	233.8	236.8	151.9	150.8	152.9
Newfoundland and Labrador	194.3	187.6	201.1	250.8	239.2	262.5	151.3	143.2	159.4
St. John's (NF1)	186.9	175.5	198.4	235.3	215.5	255.1	153.6	139.6	167.5
Eastern (NF2)	224.7	209.4	240.0	317.9	290.6	345.2	150.9	133.9	167.9
Central (NF3)	182.1	168.1	196.1	222.9	199.9	245.9	150.8	133.1	168.4
Western (NF4)	188.3	172.0	204.7	236.8	209.2	264.4	150.5	130.5	170.5
Grenfell (NF5)	168.5	129.3	207.7	190.1	128.0	252.2	149.2	98.6	199.7
Labrador (NF6)	212.0	156.6	267.5	254.0	172.3	335.7	164.5	88.9	240.0
Prince Edward Island	193.8	181.0	206.5	261.0	238.6	283.4	146.5	131.3	161.6
PEI/Urban	205.3	185.9	224.7	257.7	224.7	290.6	169.3	145.4	193.2
PEI/Rural	183.8	166.8	200.7	263.6	233.0	294.2	126.1	106.9	145.4
Nova Scotia	208.3	203.3	213.3	262.8	254.1	271.5	172.8	166.6	179.0
South-SW (NS1)	198.9	186.7	211.1	235.9	216.1	255.6	172.0	156.3	187.7
Valley (NS2)	196.1	180.2	212.0	251.2	224.1	278.4	157.0	137.6	176.4
Colch-Cumb-E.Hants (NS3)	205.1	191.0	219.3	256.2	232.3	280.0	170.6	152.9	188.2
Pictou-GASHA (NS4)	209.8	195.1	224.5	269.8	244.6	294.9	168.9	150.8	187.0
Cape Breton (NS5)	231.8	218.7	244.9	308.1	284.5	331.6	186.3	170.4	202.2
Capital (NS6)	206.8	198.1	215.6	261.2	245.6	276.8	174.4	163.8	185.0
New Brunswick	196.5	191.0	202.0	257.6	248.0	267.3	153.1	146.5	159.6
Moncton (NB1)	196.8	185.8	207.7	258.9	239.8	278.0	153.1	139.9	166.2
Sussex/Saint John (NB2)	201.2	189.9	212.5	253.6	234.0	273.3	167.1	153.2	181.1
Fredericton (NB3)	184.3	172.6	196.0	235.7	215.5	255.8	146.9	132.8	161.0
Edmundston (NB4)	199.4	178.6	220.1	288.2	249.8	326.6	134.3	111.6	157.0
Campbellton (NB5)	215.3	188.1	242.6	305.9	256.4	355.4	150.8	119.9	181.8
Bathurst (NB6)	190.3	174.0	206.7	247.2	219.3	275.1	145.2	125.8	164.5
Miramichi (NB7)	212.4	190.1	234.6	285.6	247.0	324.2	154.0	128.2	179.8

Table 142. Lung cancer deaths (ICD-9 162), by sex, crude count and confidence interval, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Crude count	95% CI		Crude count	95% CI		Crude count	95% CI	
		low	high		low	high		low	high
Canada	15,404	15,264	15,545	9,801	9,689	9,913	5,604	5,519	5,689
Newfoundland and Labrador	270	252	289	202	187	219	67	59	77
St. John's (NF1)	90	80	102	60	52	70	30	24	37
Eastern (NF2)	76	66	86	62	54	71	14	10	18
Central (NF3)	49	41	57	37	31	44	12	8	16
Western (NF4)	41	35	49	32	26	39	9	6	13
Grenfell (NF5)	6	4	10	6	4	9	--	--	2
Labrador (NF6)	7	5	11	5	3	8	2	1	5
Prince Edward Island	84	74	94	51	43	60	33	27	40
PEI/Urban	38	31	45	21	16	26	17	13	22
PEI/Rural	46	39	54	30	25	37	16	12	21
Nova Scotia	600	573	628	378	357	401	222	205	239
South-SW (NS1)	87	77	98	54	46	63	33	27	40
Valley (NS2)	49	41	57	32	26	39	17	12	22
Colch-Cumb-E.Hants (NS3)	78	69	89	51	44	60	27	22	33
Pictou-GASHA (NS4)	71	62	81	48	40	56	23	18	29
Cape Breton (NS5)	120	108	133	79	70	90	40	34	48
Capital (NS6)	195	180	212	114	102	126	82	72	92
New Brunswick	449	425	473	307	288	327	142	129	156
Moncton (NB1)	112	100	124	74	65	84	38	31	45
Sussex/Saint John (NB2)	116	104	128	73	63	83	43	36	51
Fredericton (NB3)	81	71	91	57	49	66	24	19	30
Edmundston (NB4)	40	33	48	33	27	40	7	4	10
Campbellton (NB5)	26	20	32	17	13	23	8	6	12
Bathurst (NB6)	44	37	52	32	26	39	12	9	16
Miramichi (NB7)	31	25	38	21	17	27	10	7	14

Table 143. Lung cancer deaths (ICD-9 162), by sex, crude rate per 100,000 population and confidence interval, Canada, Atlantic Provinces and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	51.9	51.4	52.4	66.7	65.9	67.5	37.4	36.8	38.0
Newfoundland and Labrador	48.1	44.9	51.5	72.4	66.8	78.4	23.9	20.8	27.4
St. John's (NF1)	48.4	42.9	54.4	65.8	56.8	76.0	31.6	25.6	38.7
Eastern (NF2)	60.7	53.2	69.0	100.1	86.5	115.3	21.8	15.9	29.3
Central (NF3)	43.0	36.4	50.3	64.8	53.5	77.7	20.8	14.7	28.5
Western (NF4)	44.6	37.3	53.0	69.2	56.3	84.1	20.1	13.6	28.7
Grenfell (NF5)	35.4	21.9	54.1	65.5	40.1	101.3	3.8	0.2	18.2
Labrador (NF6)	28.5	18.3	42.4	37.7	21.9	60.6	18.8	8.2	36.9
Prince Edward Island	61.4	54.2	69.4	75.9	64.6	88.6	47.3	38.6	57.4
PEI/Urban	59.3	49.1	71.0	66.5	51.4	84.6	52.4	39.4	68.3
PEI/Rural	63.3	53.4	74.5	84.0	68.0	102.6	42.8	31.8	56.5
Nova Scotia	64.4	61.5	67.5	82.8	78.1	87.7	46.7	43.3	50.4
South-SW (NS1)	68.2	60.3	76.9	86.2	73.7	100.2	50.6	41.3	61.4
Valley (NS2)	58.3	49.4	68.3	77.6	63.2	94.3	39.4	29.6	51.5
Colch-Cumb-E.Hants (NS3)	73.8	64.8	83.6	98.2	83.6	114.6	50.1	40.0	61.9
Pictou-GASHA (NS4)	71.0	61.9	81.0	96.5	81.7	113.3	46.1	36.2	57.9
Cape Breton (NS5)	83.9	75.6	92.9	114.5	100.6	129.7	55.0	45.8	65.5
Capital (NS6)	52.6	48.5	57.0	62.6	56.2	69.5	43.0	37.9	48.7
New Brunswick	59.6	56.5	62.9	82.1	77.0	87.6	37.4	34.0	41.1
Moncton (NB1)	61.3	55.0	68.1	81.7	71.5	93.0	41.2	34.1	49.2
Sussex/Saint John (NB2)	65.0	58.5	72.1	83.4	72.8	95.0	47.4	39.7	56.1
Fredericton (NB3)	48.8	42.9	55.2	68.6	58.9	79.5	29.0	22.8	36.3
Edmundston (NB4)	73.1	60.9	87.1	122.1	99.7	147.9	25.3	16.1	37.9
Campbellton (NB5)	77.9	61.9	96.8	106.6	80.4	138.6	49.9	33.0	72.5
Bathurst (NB6)	49.3	41.4	58.2	71.2	58.0	86.5	27.1	19.2	37.0
Miramichi (NB7)	62.0	50.3	75.6	84.5	65.6	107.1	39.1	26.7	55.3

Table 144. Lung cancer deaths (ICD-9 162), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	49.2	48.8	49.7	72.0	71.2	72.8	32.5	32.0	33.0
Newfoundland and Labrador	50.0	46.5	53.4	81.9	75.4	88.5	23.4	20.2	26.7
St. John's (NF1)	50.1	44.1	56.1	76.8	65.5	88.1	30.7	24.3	37.1
Eastern (NF2)	62.7	54.5	70.9	111.6	95.6	127.7	21.2	14.7	27.8
Central (NF3)	41.0	34.3	47.6	65.9	53.6	78.3	18.9	12.6	25.2
Western (NF4)	45.0	37.0	52.9	74.6	59.4	89.8	19.1	12.0	26.2
Grenfell (NF5)	43.1	23.5	62.7	83.0	44.1	122.0	--	--	--
Labrador (NF6)	72.9	37.6	108.2	94.4	37.8	151.0	55.7	6.7	104.7
Prince Edward Island	56.6	49.5	63.7	78.0	65.7	90.4	41.7	33.3	50.2
PEI/Urban	55.0	44.8	65.3	69.1	51.9	86.3	46.7	33.6	59.7
PEI/Rural	58.0	48.2	67.8	85.4	67.8	102.9	37.4	26.5	48.4
Nova Scotia	57.8	55.1	60.4	84.1	79.2	89.0	38.9	35.9	41.9
South-SW (NS1)	51.5	45.1	57.8	70.3	59.5	81.2	36.5	29.1	43.8
Valley (NS2)	50.1	41.9	58.3	73.9	59.1	88.7	33.2	23.9	42.5
Colch-Cumb-E.Hants (NS3)	60.5	52.7	68.2	89.4	75.3	103.5	37.5	29.2	45.9
Pictou-GASHA (NS4)	57.8	49.9	65.6	87.1	72.8	101.4	36.5	27.7	45.3
Cape Breton (NS5)	69.4	62.2	76.6	109.0	95.1	123.0	42.8	35.1	50.6
Capital (NS6)	57.1	52.5	61.8	80.2	71.6	88.8	41.9	36.6	47.2
New Brunswick	55.2	52.2	58.1	86.3	80.7	91.9	31.9	28.8	34.9
Moncton (NB1)	53.8	48.0	59.5	81.5	70.8	92.3	33.0	26.8	39.2
Sussex/Saint John (NB2)	58.6	52.4	64.8	85.8	74.4	97.2	39.7	32.7	46.7
Fredericton (NB3)	47.8	41.7	53.8	75.8	64.3	87.2	26.4	20.2	32.5
Edmundston (NB4)	67.2	55.2	79.3	131.3	105.2	157.5	20.6	11.7	29.5
Campbellton (NB5)	69.1	53.6	84.5	107.6	78.2	136.9	41.5	25.1	57.9
Bathurst (NB6)	50.0	41.4	58.6	79.2	63.3	95.1	25.0	16.7	33.3
Miramichi (NB7)	57.5	45.8	69.3	86.0	64.9	107.0	33.5	21.1	45.9

Table 145. Colorectal cancer deaths (ICD-9 153-154), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Crude count	95% CI		Crude count	95% CI		Crude count	95% CI	
		low	high		low	high		low	high
Canada	6,150	6,061	6,239	3,248	3,184	3,313	2,901	2,841	2,963
Newfoundland and Labrador	105	94	117	55	47	64	51	43	59
St. John's (NF1)	33	27	40	17	13	23	16	12	21
Eastern (NF2)	29	23	36	15	11	19	14	11	19
Central (NF3)	22	17	27	11	8	15	11	7	15
Western (NF4)	17	13	22	9	6	13	8	5	11
Grenfell (NF5)	3	1	5	1	--	3	1	--	3
Labrador (NF6)	2	1	4	1	--	3	1	--	2
Prince Edward Island	28	22	34	13	9	18	15	11	20
PEI/Urban	13	9	18	6	3	9	7	5	11
PEI/Rural	15	11	20	7	5	11	8	5	11
Nova Scotia	192	176	208	98	87	110	94	83	105
South-SW (NS1)	32	26	39	14	10	18	18	14	24
Valley (NS2)	23	18	29	11	8	16	12	8	16
Colch-Cumb-E.Hants (NS3)	24	19	30	12	9	16	12	9	17
Pictou-GASHA (NS4)	19	15	24	10	7	14	9	6	13
Cape Breton (NS5)	33	27	40	20	15	26	13	9	18
Capital (NS6)	60	52	70	31	25	38	29	24	36
New Brunswick	139	126	152	75	65	85	64	55	74
Moncton (NB1)	38	32	46	21	17	27	17	13	22
Sussex/Saint John (NB2)	33	27	40	15	11	20	17	13	23
Fredericton (NB3)	29	23	35	15	11	20	13	10	18
Edmundston (NB4)	12	9	17	6	4	10	6	4	9
Campbellton (NB5)	2	1	5	1	--	3	1	--	3
Bathurst (NB6)	16	12	21	10	7	14	6	3	9
Miramichi (NB7)	8	6	12	5	3	8	3	2	6

Table 146. Colorectal cancer deaths (ICD-9 153-154), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	20.7	20.4	21.0	22.1	21.7	22.6	19.4	19.0	19.8
Newfoundland and Labrador	18.8	16.8	20.9	19.6	16.7	22.7	18.0	15.3	21.1
St. John's (NF1)	17.9	14.6	21.6	18.9	14.3	24.6	16.9	12.6	22.1
Eastern (NF2)	23.3	18.8	28.6	23.7	17.4	31.5	22.9	16.8	30.5
Central (NF3)	19.1	14.9	24.2	19.3	13.5	26.7	19.0	13.2	26.5
Western (NF4)	18.4	13.8	23.9	20.2	13.7	28.7	16.6	10.7	24.4
Grenfell (NF5)	14.9	6.9	28.2	14.6	4.7	34.8	15.2	4.9	36.4
Labrador (NF6)	6.5	2.4	14.3	7.5	1.9	20.3	5.4	0.9	17.4
Prince Edward Island	20.6	16.5	25.3	19.3	13.9	26.2	21.7	16.0	28.8
PEI/Urban	20.5	14.8	27.7	18.2	11.0	28.5	22.6	14.5	33.6
PEI/Rural	20.6	15.2	27.4	20.3	13.0	30.2	21.0	13.6	30.9
Nova Scotia	20.6	18.9	22.3	21.5	19.1	24.0	19.7	17.5	22.2
South-SW (NS1)	25.1	20.4	30.5	21.7	15.8	29.1	28.4	21.6	36.7
Valley (NS2)	27.6	21.6	34.6	27.5	19.3	37.9	27.6	19.5	37.9
Colch-Cumb-E.Hants (NS3)	22.9	18.1	28.6	23.0	16.3	31.4	22.9	16.3	31.2
Pictou-GASHA (NS4)	19.0	14.5	24.4	20.3	13.9	28.5	17.8	12.0	25.5
Cape Breton (NS5)	23.1	18.9	28.0	28.9	22.2	36.9	17.7	12.8	24.0
Capital (NS6)	16.2	14.0	18.7	17.1	13.9	20.8	15.5	12.5	18.9
New Brunswick	18.4	16.7	20.2	20.0	17.5	22.7	16.9	14.6	19.4
Moncton (NB1)	21.0	17.4	25.1	23.6	18.3	29.9	18.4	13.9	24.0
Sussex/Saint John (NB2)	18.4	15.0	22.3	17.6	13.0	23.2	19.1	14.4	24.9
Fredericton (NB3)	17.3	13.9	21.3	18.6	13.7	24.5	16.1	11.7	21.7
Edmundston (NB4)	22.5	16.1	30.7	23.4	14.5	35.8	21.7	13.2	33.5
Campbellton (NB5)	7.1	3.1	13.9	6.1	1.6	16.5	8.0	2.6	19.1
Bathurst (NB6)	17.9	13.4	23.5	23.0	15.9	32.2	12.8	7.7	20.0
Miramichi (NB7)	16.7	11.0	24.2	19.8	11.5	31.9	13.5	6.8	23.9

Table 147. Colorectal cancer deaths (ICD-9 153-154), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	19.4	19.1	19.7	24.4	23.9	24.9	15.7	15.4	16
Newfoundland and Labrador	19.4	17.3	21.6	22.7	19.2	26.2	16.9	14.2	19.7
St. John's (NF1)	18.2	14.6	21.8	22.0	15.9	28.1	15.7	11.2	20.2
Eastern (NF2)	23.1	18.2	27.9	27.4	19.2	35.6	20.1	14.0	26.1
Central (NF3)	18.6	14.1	23.2	20.0	13.1	26.9	17.3	11.2	23.3
Western (NF4)	19.4	14.0	24.8	22.7	14.2	31.2	15.9	9.4	22.4
Grenfell (NF5)	16.7	5.0	28.4	17.7	0.1	35.3	15.7	0.2	31.1
Labrador (NF6)	--	--	--	--	--	--	--	--	--
Prince Edward Island	17.5	13.7	21.2	19.3	13.3	25.4	15.9	11.1	20.8
PEI/Urban	17.3	11.8	22.8	17.7	9.3	26.1	17.2	9.7	24.8
PEI/Rural	17.5	12.3	22.7	20.5	11.9	29.1	14.8	8.6	21.1
Nova Scotia	17.7	16.2	19.1	22.0	19.4	24.5	14.6	12.9	16.4
South-SW (NS1)	17.5	13.9	21.0	17.3	12.0	22.7	17.6	12.7	22.5
Valley (NS2)	22.3	17.0	27.7	26.1	17.3	34.9	18.6	12.2	25.0
Colch-Cumb-E.Hants (NS3)	17.5	13.5	21.6	20.8	14.0	27.6	14.6	9.7	19.4
Pictou-GASHA (NS4)	13.8	10.1	17.4	18.4	11.8	25.1	11.1	6.7	15.5
Cape Breton (NS5)	18.3	14.7	22.0	28.5	21.2	35.8	12.2	8.2	16.1
Capital (NS6)	17.1	14.6	19.7	21.7	17.2	26.1	14.2	11.2	17.2
New Brunswick	16.4	14.8	18.0	21.0	18.2	23.8	13.0	11.1	14.8
Moncton (NB1)	17.6	14.4	20.8	23.4	17.6	29.2	14.0	10.0	17.9
Sussex/Saint John (NB2)	15.7	12.5	18.8	18.3	13.0	23.6	13.2	9.4	16.9
Fredericton (NB3)	16.5	13.0	20.0	20.7	14.7	26.8	13.3	9.1	17.6
Edmundston (NB4)	20.0	13.6	26.5	24.2	13.2	35.1	16.5	8.8	24.1
Campbellton (NB5)	6.1	1.6	10.6	--	--	--	--	--	--
Bathurst (NB6)	17.8	12.8	22.9	25.9	16.8	35.1	11.4	5.9	17
Miramichi (NB7)	15.3	9.3	21.3	19.9	9.8	30.0	10.1	3.7	16.6

Table 148. Prostate cancer deaths (ICD-9 185), crude count, crude rate per 100,000 population, age-standardized rate per 100,000 population and confidence intervals, Canada, Atlantic Provinces, and Health Districts, 1996

	Crude count			Crude rate per 100,000 population			Age-standardized rate per 100,000 population		
	Crude count	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	3,657	3,589	3,726	24.9	24.4	25.4	29.7	29.1	30.2
Newfoundland and Labrador	65	56	74	23.1	20.1	26.6	29.8	25.5	34.1
St. John's (NF1)	18	14	24	20.0	15.2	25.8	25.5	18.6	32.4
Eastern (NF2)	19	15	25	31.2	23.9	40	38.7	28.5	48.9
Central (NF3)	14	10	19	24.5	17.9	32.8	28.2	19.5	36.9
Western (NF4)	12	8	16	25.2	17.8	34.7	32.4	21.3	43.4
Grenfell (NF5)	1	0	2	7.3	1.3	23.6
Labrador (NF6)	1	0	2	5.0	0.9	16.3
Prince Edward Island	24	19	30	35.2	27.7	44.1	36.0	27.6	44.4
PEI/Urban	10	7	14	32.2	22.1	45.3	33.0	21.2	44.8
PEI/Rural	14	10	18	37.8	27.5	50.8	38.5	26.7	50.3
Nova Scotia	142	129	156	31.2	28.3	34.2	33.3	30.1	36.5
South-SW (NS1)	24	19	30	37.6	29.5	47.1	29.7	22.7	36.6
Valley (NS2)	18	13	23	42.9	32.4	55.6	41.8	30.5	53.0
Colch-Cumb-E.Hants (NS3)	18	13	23	33.8	25.6	43.8	31.4	22.9	39.8
Pictou-GASHA (NS4)	16	12	21	31.7	23.6	41.8	27.5	19.6	35.4
Cape Breton (NS5)	24	19	30	34.6	27.3	43.3	35.3	27.0	43.6
Capital (NS6)	44	37	52	24.0	20.2	28.4	35.1	28.9	41.2
New Brunswick	100	89	111	26.7	23.8	29.8	29.9	26.5	33.3
Moncton (NB1)	31	25	37	33.9	27.5	41.3	34.7	27.6	41.9
Sussex/Saint John (NB2)	23	18	29	26.4	20.7	33.2	28.2	21.4	34.9
Fredericton (NB3)	17	13	22	20.6	15.5	26.8	25.3	18.3	32.3
Edmundston (NB4)	5	3	8	19.7	11.7	31.3	22.5	11.3	33.7
Campbellton (NB5)	4	2	6	22.5	11.9	39.1	25.0	10.2	39.7
Bathurst (NB6)	14	11	19	31.9	23.4	42.5	38.8	27.3	50.4
Miramichi (NB7)	6	3	9	22.4	13.5	35.1	25.6	13.3	37.9

Table 149. Female breast cancer deaths (ICD-9 174), crude count, crude rate per 100,000 population, age-standardized rate per 100,000 population and confidence intervals, Canada, Atlantic Provinces, and Health Districts, 1996

	Crude count			Crude rate per 100,000 population			Age-standardized rate per 100,000 population		
	Crude count	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	4,980	4,900	5,060	33.2	32.7	33.8	28.3	27.9	28.8
Newfoundland and Labrador	80	70	91	28.4	25.0	32.2	27.0	23.6	30.5
St. John's (NF1)	31	25	37	32.3	26.2	39.4	29.9	23.7	36.1
Eastern (NF2)	13	10	18	21.3	15.4	28.7	20.3	13.9	26.7
Central (NF3)	17	13	22	30.3	22.8	39.5	27.8	20.1	35.4
Western (NF4)	17	13	23	37.4	28.2	48.7	34.7	25.2	44.2
Grenfell (NF5)	1	--	3	15.2	4.9	36.4	--	--	--
Labrador (NF6)	--	--	2	2.7	0.1	12.8	--	--	--
Prince Edward Island	21	16	26	30.0	23.2	38.1	23.8	17.7	30.0
PEI/Urban	11	8	15	33.9	23.7	47.0	27.2	17.5	37.0
PEI/Rural	10	7	14	26.4	18.0	37.4	20.8	13.0	28.6
Nova Scotia	185	170	201	38.9	35.8	42.3	30.9	28.2	33.5
South-SW (NS1)	29	23	35	44.4	35.7	54.6	29.0	22.6	35.5
Valley (NS2)	15	11	19	34.7	25.5	46.1	27.3	18.9	35.6
Colch-Cumb-E.Hants (NS3)	20	15	26	37.1	28.5	47.4	28.2	20.9	35.6
Pictou-GASHA (NS4)	20	16	26	40.2	31.0	51.2	29.6	21.8	37.3
Cape Breton (NS5)	33	27	40	44.5	36.4	54.0	33.9	27.0	40.8
Capital (NS6)	68	59	78	36.0	31.3	41.2	32.9	28.3	37.5
New Brunswick	135	122	148	35.5	32.2	39.1	29.3	26.4	32.2
Moncton (NB1)	36	30	43	39.0	32.1	46.9	31.1	25.1	37.2
Sussex/Saint John (NB2)	35	29	42	38.6	31.7	46.5	30.7	24.7	36.7
Fredericton (NB3)	26	21	32	31.4	25.0	39.0	26.6	20.6	32.6
Edmundston (NB4)	9	6	12	31.3	20.9	45.2	26.6	16.3	37.0
Campbellton (NB5)	6	4	9	36.0	22.0	55.6	29.7	15.9	43.5
Bathurst (NB6)	15	11	20	33.8	25.0	44.9	29.5	20.8	38.2
Miramichi (NB7)	8	5	12	32.3	21.2	47.3	28.1	16.7	39.5

Table 150. Age standardized respiratory disease death rate (ICD-9 460-519), Canada, Atlantic Provinces, and Health Districts, 1996

	All respiratory disease deaths						Pneumonia and influenza disease deaths						Bronchitis, emphysema and asthma disease deaths						All other respiratory disease deaths					
	Both sexes		Male		Female		Both sexes		Male		Female		Both sexes		Male		Female		Both sexes		Male		Female	
	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province
Canada																								
Newfoundland and Labrador	✓		×		✓		✓		×		✓		✓		✓		✓		×		×		✓	
St. John's (NF1)	✓	✓	✓	✓	✓	×	✓	✓	×	×	✓	✓	✓	×	✓	×	✓	×	✓	✓	✓	✓	×	×
Eastern (NF2)	×	×	×	×	✓	×	×	×	×	×	✓	×	✓	✓	✓	✓	✓	×	×	×	×	×	✓	×
Central (NF3)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	-	-	-	-	✓	✓	×	×	✓	✓	
Western (NF4)	×	×	×	×	✓	×	×	×	×	×	×	×	×	✓	×	✓	×	×	×	×	×	✓	×	
Grenfell (NF5)	✓	✓	✓	✓	✓	✓	✓	✓	-	-	-	-	-	-	-	-	-	✓	✓	✓	✓	-	-	
Labrador (NF6)	×	×	×	×	×	×	×	×	×	-	-	-	-	-	-	-	-	×	×	×	×	×	×	
Prince Edward Island	×		×		×		×		×		×		×		×		×		✓		✓		✓	
PEI/Urban	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	s	×	×	✓	×	×	×	
PEI/Rural	✓	✓	×	✓	✓	✓	×	✓	×	✓	✓	✓	×	✓	×	✓	×	s	✓	✓	✓	✓	✓	✓
Nova Scotia	×		×		×		×		×		×		✓		×		✓		×		×		×	
South-SW (NS1)	×	✓	✓	✓	✓	✓	×	✓	×	✓	×	✓	✓	×	s	✓	✓	✓	✓	✓	✓	✓	✓	✓
Valley (NS2)	×	×	×	×	×	✓	×	×	×	×	×	×	×	×	×	✓	✓	×	✓		✓	✓	✓	
Colch-Cumb-E.Hants (NS3)	×	×	×	✓	×	×	×	✓	×	✓	×	✓	✓	✓	✓	✓	✓	✓	×	×	×	✓	×	
Pictou-GASHA (NS4)	×	×	×	✓	×	×	×	×	×	×	×	✓	✓	✓	✓	✓	✓	✓	×	✓	×	✓	×	
Cape Breton (NS5)	×	×	×	×	×	×	✓	✓	✓	✓	×	✓	×	×	×	×	×	×	×	×	×	×	×	
Capital (NS6)	×	✓	×	×	×	×	×	×	×	×	✓	✓	✓	✓	✓	✓	✓	✓	×	×	×	×	×	
New Brunswick	✓		✓		✓		✓		✓		✓		✓		✓		✓		×		×		×	
Moncton (NB1)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	×	×	✓	✓	✓	✓	✓	✓
Sussex/Saint John (NB2)	×	×	×	×	×	×	✓	×	×	×	✓	×	✓	×	×	×	✓	✓	×	×	×	×	×	
Fredericton (NB3)	×	×	×	×	×	×	✓	×	×	×	×	×	✓	×	✓	×	✓	✓	×	×	×	×	×	
Edmundston (NB4)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	×	×	×	×	✓	✓	×	×	×	×	×	
Campbellton (NB5)	✓	✓	✓	✓	×	×	✓	✓	✓	✓	✓	✓	✓	-	-	-	-	×	×	×	✓	×	×	
Bathurst (NB6)	✓	✓	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	×	×	×	×	✓	✓	×	✓	×	×	✓	
Miramichi (NB7)	✓	✓	✓	✓	×	×	✓	✓	✓	✓	✓	✓	×	✓	✓	×	✓	-	-	×	✓	✓	✓	

× - less favourable rate than the province or nation
s - same rate as the province or nation

✓ - more favourable rate than the province or nation
“-” - no data available for comparison

Table 151. All respiratory disease deaths (ICD-9 460-519), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Crude count	95% CI		Crude count	95% CI		Crude count	95% CI	
		low	high		low	high		low	high
Canada	19,348	19,191	19,506	10,369	10,255	10,485	8,979	8,872	9,086
Newfoundland and Labrador	303	284	324	191	176	207	112	101	125
St. John's (NF1)	102	91	114	59	50	68	44	37	52
Eastern (NF2)	77	67	87	49	42	58	27	22	34
Central (NF3)	52	45	61	37	31	45	15	11	20
Western (NF4)	57	49	66	36	30	43	21	16	27
Grenfell (NF5)	5	3	8	4	2	6	2	1	4
Labrador (NF6)	10	7	14	6	4	9	4	2	6
Prince Edward Island	119	107	132	64	55	74	55	47	64
PEI/Urban	64	55	74	30	24	37	34	28	41
PEI/Rural	55	47	64	34	28	41	21	16	27
Nova Scotia	814	782	846	440	417	465	373	352	396
South-SW (NS1)	112	100	124	63	54	72	49	42	57
Valley (NS2)	84	74	94	47	40	56	36	30	44
Colch-Cumb-E.Hants (NS3)	106	95	118	54	46	63	52	44	61
Pictou-GASHA (NS4)	106	95	118	58	50	67	48	41	57
Cape Breton (NS5)	141	128	155	77	67	87	64	55	74
Capital (NS6)	266	248	285	142	129	156	124	112	137
New Brunswick	509	484	535	274	256	294	235	218	252
Moncton (NB1)	105	94	117	55	47	64	50	43	59
Sussex/Saint John (NB2)	148	135	162	77	68	88	71	62	81
Fredericton (NB3)	122	110	135	67	58	76	55	47	64
Edmundston (NB4)	33	27	40	19	15	25	14	10	19
Campbellton (NB5)	21	17	27	11	8	15	10	7	14
Bathurst (NB6)	50	42	58	32	26	39	18	14	24
Miramichi (NB7)	29	24	36	14	10	18	16	12	21

Table 152. All respiratory disease deaths (ICD-9 460-519), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	65.2	64.7	65.7	70.6	69.8	71.4	59.9	59.2	60.7
Newfoundland and Labrador	54.1	50.7	57.7	68.4	62.9	74.1	39.9	35.8	44.4
St. John's (NF1)	54.9	49.0	61.2	64.0	55.1	74.0	46.0	38.6	54.4
Eastern (NF2)	61.5	54.0	69.9	79.7	67.6	93.3	43.6	34.9	53.9
Central (NF3)	46.2	39.4	53.8	65.3	54.0	78.3	26.7	19.7	35.4
Western (NF4)	61.6	52.8	71.3	77.8	64.1	93.5	45.3	35.1	57.6
Grenfell (NF5)	29.8	17.6	47.2	40.1	21.1	69.4	19.1	7.0	41.9
Labrador (NF6)	37.6	25.7	53.3	45.2	27.6	69.9	29.5	15.5	51.1
Prince Edward Island	87.4	78.7	96.8	95.3	82.5	109.4	79.7	68.2	92.6
PEI/Urban	100.8	87.3	115.8	96.5	78.1	118.0	104.8	85.9	126.7
PEI/Rural	75.7	64.8	87.9	94.1	77.1	113.8	57.4	44.5	73.0
Nova Scotia	87.4	84.0	90.9	96.4	91.3	101.7	78.7	74.2	83.4
South-SW (NS1)	87.5	78.5	97.3	99.4	86.0	114.4	75.9	64.4	89.0
Valley (NS2)	100.2	88.4	113.2	114.8	97.0	134.9	86.0	70.9	103.3
Colch-Cumb-E.Hants (NS3)	99.8	89.3	111.2	103.3	88.3	120.1	96.4	82.2	112.4
Pictou-GASHA (NS4)	106.0	94.8	118.1	116.8	100.3	135.2	95.5	80.8	112.0
Cape Breton (NS5)	98.6	89.5	108.4	110.6	97.0	125.6	87.3	75.6	100.3
Capital (NS6)	71.5	66.7	76.6	78.2	71.0	85.9	65.2	58.8	72.1
New Brunswick	67.6	64.3	71.0	73.4	68.5	78.5	61.9	57.4	66.6
Moncton (NB1)	57.4	51.3	64.0	60.4	51.7	70.2	54.5	46.3	63.7
Sussex/Saint John (NB2)	83.2	75.7	91.2	88.7	77.8	100.7	77.9	68.0	89.0
Fredericton (NB3)	73.8	66.5	81.6	80.7	70.1	92.5	66.8	57.2	77.6
Edmundston (NB4)	60.9	49.8	73.8	71.5	54.8	91.7	50.6	36.9	67.7
Campbellton (NB5)	64.7	50.3	82.1	67.6	47.3	93.8	61.9	42.8	86.7
Bathurst (NB6)	56.0	47.6	65.5	70.5	57.3	85.7	41.4	31.5	53.4
Miramichi (NB7)	58.7	47.3	71.9	54.1	39.3	72.7	63.3	47.1	83.4

Table 153. All respiratory disease deaths (ICD-9 460-519), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	59.8	59.3	60.3	86.2	85.2	87.1	44.3	43.7	44.8
Newfoundland and Labrador	56.9	53.2	60.6	90.3	82.8	97.9	36.0	32.1	39.8
St. John's (NF1)	54.8	48.7	61.0	85.1	72.2	98.0	38.3	31.6	44.9
Eastern (NF2)	61.4	53.5	69.4	103.8	86.8	120.9	37.2	29.1	45.3
Central (NF3)	45.2	38.1	52.3	75.0	60.9	89.1	23.4	16.5	30.3
Western (NF4)	67.6	57.4	77.7	102.2	82.3	122.1	43.1	32.5	53.7
Grenfell (NF5)	43.4	21.6	65.2	60.5	22.2	98.7	27.6	3.5	51.8
Labrador (NF6)	121.3	73.5	169.2	146.7	73.5	219.8	92.9	33.9	151.9
Prince Edward Island	66.5	59.6	73.5	98.8	84.9	112.7	47.5	39.9	55.2
PEI/Urban	76.6	65.6	87.5	103.6	82.4	124.9	62.7	49.9	75.6
PEI/Rural	57.8	48.9	66.8	94.7	76.4	113.1	34.2	25.3	43.1
Nova Scotia	71.0	68.2	73.9	105.2	99.5	110.9	51.3	48.2	54.4
South-SW (NS1)	55.7	49.7	61.7	81.0	69.3	92.6	39.3	32.7	45.9
Valley (NS2)	72.9	63.9	82.0	112.3	93.9	130.6	50.1	40.3	59.8
Colch-Cumb-E.Hants (NS3)	70.7	62.9	78.5	96.3	81.5	111.1	54.2	45.4	62.9
Pictou-GASHA (NS4)	72.2	64.2	80.3	104.7	89.1	120.3	52.2	43.2	61.1
Cape Breton (NS5)	76.3	69.0	83.7	115.7	100.5	131.0	54.2	46.3	62.1
Capital (NS6)	74.7	69.5	79.9	115.6	104.4	126.8	54.2	48.6	59.8
New Brunswick	57.4	54.5	60.3	82.2	76.5	87.8	43.1	39.9	46.4
Moncton (NB1)	45.4	40.4	50.5	63.5	53.8	73.3	35.3	29.5	41.1
Sussex/Saint John (NB2)	66.8	60.5	73.1	96.6	84.0	109.2	51.9	44.7	59.2
Fredericton (NB3)	67.2	60.3	74.1	96.4	83.0	109.7	48.6	41.1	56.2
Edmundston (NB4)	55.0	44.2	65.8	82.0	60.7	103.3	37.7	26.3	49.2
Campbellton (NB5)	54.4	41.1	67.7	75.0	49.5	100.4	44.8	28.8	60.8
Bathurst (NB6)	52.2	43.9	60.6	84.5	67.6	101.3	31.3	22.9	39.7
Miramichi (NB7)	51.1	40.4	61.8	57.4	39.7	75.1	46.2	32.7	59.6

Table 154. Pneumonia and influenza deaths (ICD-9 480-487), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Crude count	95% CI		Crude count	95% CI		Crude count	95% CI	
		low	high		low	high		low	high
Canada	7,627	7,529	7,727	3,585	3,518	3,654	4,042	3,971	4,114
Newfoundland and Labrador	112	100	124	66	57	76	46	38	54
St. John's (NF1)	38	32	46	22	17	27	17	12	22
Eastern (NF2)	30	24	37	19	15	25	11	7	15
Central (NF3)	17	13	22	10	7	14	7	5	11
Western (NF4)	22	17	28	13	9	17	9	6	13
Grenfell (NF5)	2	1	4	1	--	3	1	--	2
Labrador (NF6)	3	1	5	2	1	4	1	--	3
Prince Edward Island	61	52	70	30	24	37	31	25	37
PEI/Urban	32	26	39	14	10	18	18	14	23
PEI/Rural	29	23	36	16	12	21	13	9	17
Nova Scotia	344	324	366	158	145	173	186	171	202
South-SW (NS1)	55	47	64	26	21	32	29	23	35
Valley (NS2)	41	34	48	19	15	24	22	17	27
Colch-Cumb-E.Hants (NS3)	43	36	51	20	15	25	23	18	29
Pictou-GASHA (NS4)	51	43	60	23	18	29	28	22	34
Cape Breton (NS5)	44	37	52	15	11	19	29	23	36
Capital (NS6)	112	100	124	56	48	65	56	48	65
New Brunswick	158	144	173	69	60	79	89	79	101
Moncton (NB1)	33	27	40	16	12	21	18	13	23
Sussex/Saint John (NB2)	49	41	57	23	18	29	26	21	32
Fredericton (NB3)	42	35	49	16	12	21	26	20	32
Edmundston (NB4)	6	4	9	2	1	5	4	2	6
Campbellton (NB5)	6	4	10	3	1	5	3	2	6
Bathurst (NB6)	14	10	19	7	4	10	7	4	10
Miramichi (NB7)	8	5	12	2	1	4	6	4	9

Table 155. Pneumonia and influenza deaths (ICD-9 480-487), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	25.7	25.4	26.0	24.4	23.9	24.9	27.0	26.5	27.5
Newfoundland and Labrador	19.9	17.9	22.1	23.6	20.5	27.1	16.2	13.7	19.1
St. John's (NF1)	20.5	17.0	24.6	23.6	18.4	29.9	17.6	13.2	22.9
Eastern (NF2)	24.1	19.5	29.5	31.2	23.9	40.0	17.0	11.8	23.7
Central (NF3)	15.0	11.3	19.6	16.9	11.5	24.0	13.1	8.4	19.4
Western (NF4)	23.8	18.5	30.0	27.4	19.6	37.2	20.1	13.6	28.7
Grenfell (NF5)	9.3	3.4	20.5	10.9	2.8	29.4	7.6	1.3	24.7
Labrador (NF6)	10.4	4.8	19.6	12.6	4.6	27.6	8.0	2.1	21.6
Prince Edward Island	44.5	38.4	51.4	44.6	36.1	54.6	44.4	36.0	54.2
PEI/Urban	49.9	40.6	60.7	44.0	32.0	59.0	55.5	42.1	71.8
PEI/Rural	39.9	32.1	49.0	45.2	33.8	59.3	34.6	24.9	47.0
Nova Scotia	37.0	34.8	39.3	34.7	31.6	37.9	39.2	36.1	42.6
South-SW (NS1)	42.9	36.7	49.8	41.3	32.8	51.2	44.4	35.7	54.6
Valley (NS2)	48.7	40.6	57.9	46.1	35.2	59.2	51.3	39.9	64.9
Colch-Cumb-E.Hants (NS3)	40.2	33.6	47.6	37.6	28.9	48.2	42.6	33.4	53.6
Pictou-GASHA (NS4)	51.0	43.4	59.6	46.6	36.5	58.6	55.3	44.4	68.1
Cape Breton (NS5)	30.6	25.7	36.2	21.2	15.6	28.1	39.5	31.9	48.5
Capital (NS6)	30.1	27.0	33.4	30.8	26.4	35.8	29.3	25.1	34.0
New Brunswick	21.0	19.2	22.9	18.4	16.0	21.0	23.6	20.9	26.5
Moncton (NB1)	18.2	14.9	22.1	17.3	12.9	22.8	19.1	14.5	24.8
Sussex/Saint John (NB2)	27.4	23.2	32.1	26.0	20.3	32.7	28.7	22.8	35.6
Fredericton (NB3)	25.2	21.1	29.9	19.4	14.4	25.4	31.0	24.6	38.5
Edmundston (NB4)	11.0	6.7	17.0	8.6	3.8	17.0	13.2	7.0	22.9
Campbellton (NB5)	19.2	11.9	29.4	18.4	9.0	33.7	20.0	10.2	35.5
Bathurst (NB6)	15.7	11.4	21.0	15.6	9.9	23.4	15.8	10.0	23.7
Miramichi (NB7)	16.0	10.5	23.4	7.9	3.2	16.4	24.3	14.8	37.5

Table 156. Pneumonia and influenza deaths (ICD-9 480-487), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	23.3	23.0	23.6	31.1	30.5	31.7	18.8	18.5	19.2
Newfoundland and Labrador	21.2	18.9	23.5	33.4	28.6	38.2	14.0	11.7	16.4
St. John's (NF1)	20.3	16.6	24.1	33.6	25.1	42.1	13.5	9.7	17.3
Eastern (NF2)	24.3	19.3	29.3	44.7	32.9	56.5	14.1	9.2	19.0
Central (NF3)	14.9	10.8	18.9	20.8	13.1	28.6	10.9	6.4	15.5
Western (NF4)	26.4	20.0	32.8	36.4	24.2	48.5	18.9	11.9	25.9
Grenfell (NF5)	15.0	1.3	28.6	--	--	--	--	--	--
Labrador (NF6)	37.2	9.3	65.1	43.0	4.3	81.7	--	--	--
Prince Edward Island	32.1	27.4	36.8	48.1	38.2	58.0	23.6	18.6	28.7
PEI/Urban	36.3	28.9	43.7	48.9	34.0	63.8	30.9	22.2	39.6
PEI/Rural	28.4	22.4	34.4	47.4	34.2	60.7	17.2	11.7	22.7
Nova Scotia	29.1	27.3	30.9	39.3	35.7	42.9	23.6	21.6	25.6
South-SW (NS1)	26.1	22.1	30.2	33.7	26.1	41.3	20.9	16.3	25.5
Valley (NS2)	33.8	27.8	39.8	45.5	33.7	57.3	27.0	20.2	33.7
Colch-Cumb-E.Hants (NS3)	27.4	22.6	32.2	36.3	27.0	45.6	22.7	17.1	28.3
Pictou-GASHA (NS4)	33.4	28.1	38.7	42.8	32.6	52.9	27.7	21.6	33.8
Cape Breton (NS5)	23.2	19.2	27.2	23.2	16.1	30.3	23.2	18.2	28.3
Capital (NS6)	30.7	27.4	33.9	48.5	40.9	56.0	22.5	19.1	26.0
New Brunswick	17.3	15.7	18.8	21.3	18.3	24.2	14.9	13.1	16.8
Moncton (NB1)	13.8	11.1	16.5	19.2	13.7	24.7	11.0	8.0	14.0
Sussex/Saint John (NB2)	21.3	17.8	24.7	28.6	21.7	35.6	17.5	13.5	21.5
Fredericton (NB3)	22.2	18.3	26.1	24.2	17.3	31.1	20.5	15.9	25.2
Edmundston (NB4)	10.1	5.4	14.8	9.2	2.4	16.0	9.4	3.8	15.0
Campbellton (NB5)	15.8	8.7	22.9	20.1	6.9	33.3	12.7	4.7	20.7
Bathurst (NB6)	14.3	9.9	18.6	19.4	11.1	27.6	11.3	6.4	16.2
Miramichi (NB7)	13.5	8.1	18.8	8.7	1.6	15.9	16.3	8.6	24.0

Table 157. Bronchitis, emphysema and asthma deaths (ICD-9 490-493), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Crude count	95% CI		Crude count	95% CI		Crude count	95% CI	
		low	high		low	high		low	high
Canada	1,953	1,903	2,003	1,093	1,056	1,131	860	827	893
Newfoundland and Labrador	22	17	27	14	10	19	8	5	11
St. John's (NF1)	9	6	13	5	3	8	3	2	6
Eastern (NF2)	5	3	8	3	1	5	2	1	4
Central (NF3)	1	--	3	1	--	3	--	--	2
Western (NF4)	5	3	8	4	2	6	2	1	4
Grenfell (NF5)	--	--	2	--	--	2	--	--	--
Labrador (NF6)	1	--	3	1	--	2	--	--	2
Prince Edward Island	12	9	16	8	5	11	4	2	7
PEI/Urban	7	4	10	4	2	7	2	1	5
PEI/Rural	5	3	8	3	2	6	2	1	4
Nova Scotia	62	54	72	38	32	46	24	19	30
South-SW (NS1)	9	6	13	7	4	10	3	1	5
Valley (NS2)	7	4	10	5	3	8	2	1	4
Colch-Cumb-E.Hants (NS3)	8	5	11	5	3	8	3	1	5
Pictou-GASHA (NS4)	7	5	11	5	3	8	3	1	5
Cape Breton (NS5)	16	12	21	10	7	14	7	4	10
Capital (NS6)	15	11	20	8	5	11	7	5	11
New Brunswick	47	40	56	28	22	34	20	15	25
Moncton (NB1)	11	8	16	4	2	7	7	4	10
Sussex/Saint John (NB2)	12	8	16	8	5	11	4	2	7
Fredericton (NB3)	10	7	14	6	3	9	4	2	7
Edmundston (NB4)	4	2	7	3	1	5	1	--	3
Campbellton (NB5)	1	--	3	1	--	2	1	--	2
Bathurst (NB6)	6	4	10	5	3	8	2	1	4
Miramichi (NB7)	2	1	5	2	1	4	1	--	2

Table 158. Bronchitis, emphysema and asthma deaths (ICD-9 490-493), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	6.6	6.4	6.8	7.4	7.2	7.7	5.7	5.5	6.0
Newfoundland and Labrador	3.9	3.0	4.9	5.0	3.7	6.7	2.7	1.8	4.0
St. John's (NF1)	4.6	3.1	6.7	5.8	3.4	9.2	3.5	1.8	6.2
Eastern (NF2)	4.0	2.3	6.5	4.8	2.4	8.9	3.2	1.3	6.6
Central (NF3)	1.2	0.4	2.8	1.8	0.5	4.7	0.6	--	2.8
Western (NF4)	5.8	3.4	9.1	7.9	4.2	13.7	3.6	1.3	7.9
Grenfell (NF5)	1.9	0.1	8.9	3.6	0.2	17.4	--	--	--
Labrador (NF6)	3.9	1.0	10.5	5.0	0.9	16.3	2.7	0.1	12.8
Prince Edward Island	8.8	6.3	12.1	11.4	7.4	16.8	6.3	3.5	10.4
PEI/Urban	10.5	6.6	15.9	13.9	7.8	23.2	7.2	3.2	14.2
PEI/Rural	7.3	4.3	11.6	9.2	4.7	16.4	5.5	2.2	11.3
Nova Scotia	6.7	5.8	7.7	8.4	7.0	10.0	5.1	4.0	6.3
South-SW (NS1)	7.3	5.0	10.4	10.6	6.6	16.0	4.1	1.9	7.8
Valley (NS2)	8.0	5.0	12.1	12.1	7.0	19.5	3.9	1.5	8.7
Colch-Cumb-E.Hants (NS3)	7.2	4.7	10.6	8.9	5.1	14.6	5.6	2.7	10.2
Pictou-GASHA (NS4)	7.3	4.7	10.9	9.5	5.4	15.4	5.3	2.4	10.0
Cape Breton (NS5)	11.5	8.6	15.0	14.0	9.5	19.8	9.1	5.7	13.8
Capital (NS6)	4.0	3.0	5.4	4.2	2.7	6.2	3.9	2.5	5.7
New Brunswick	6.3	5.3	7.4	7.4	5.9	9.1	5.2	4.0	6.6
Moncton (NB1)	6.2	4.4	8.6	4.8	2.7	8.0	7.6	4.8	11.4
Sussex/Saint John (NB2)	6.6	4.6	9.0	8.8	5.7	13.0	4.4	2.4	7.5
Fredericton (NB3)	6.0	4.2	8.5	6.9	4.1	10.7	5.2	2.9	8.7
Edmundston (NB4)	7.9	4.4	13.2	11.1	5.4	20.3	4.8	1.5	11.5
Campbellton (NB5)	4.0	1.3	9.7	4.1	0.7	13.3	4.0	0.7	13.0
Bathurst (NB6)	7.1	4.4	10.9	10.4	5.9	17.0	3.8	1.4	8.3
Miramichi (NB7)	4.7	2.0	9.2	6.6	2.4	14.5	2.7	0.5	8.8

Table 159. Bronchitis, emphysema and asthma deaths (ICD-9 490-493), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	6.1	6.0	6.3	8.7	8.4	9.0	4.5	4.4	4.7
Newfoundland and Labrador	4.0	3.0	4.9	5.8	4.1	7.6	2.6	1.5	3.6
St. John's (NF1)	4.6	2.8	6.4	7.0	3.5	10.4	3.2	1.2	5.1
Eastern (NF2)	3.9	1.9	6.0	5.3	1.8	8.8	3.0	0.6	5.5
Central (NF3)	1.1	--	2.2	--	--	--	--	--	--
Western (NF4)	6.2	3.2	9.3	9.4	3.8	15.0	3.6	0.4	6.7
Grenfell (NF5)	--	--	--	--	--	--	--	--	--
Labrador (NF6)	--	--	--	--	--	--	--	--	--
Prince Edward Island	7.1	4.7	9.4	11.4	6.7	16.1	4.7	2.0	7.4
PEI/Urban	8.1	4.5	11.7	14.4	6.5	22.3	4.7	1.0	8.3
PEI/Rural	6.2	3.1	9.2	8.8	3.4	14.3	4.7	0.9	8.6
Nova Scotia	5.7	4.9	6.5	9.0	7.3	10.6	3.7	2.8	4.6
South-SW (NS1)	5.0	3.1	6.9	9.0	5.0	13.0	2.6	0.7	4.5
Valley (NS2)	6.2	3.4	8.9	11.6	5.7	17.5	2.8	0.2	5.3
Colch-Cumb-E.Hants (NS3)	5.3	3.1	7.5	8.0	3.8	12.1	3.5	1.2	5.9
Pictou-GASHA (NS4)	5.3	3.0	7.5	8.6	4.0	13.1	3.3	0.8	5.7
Cape Breton (NS5)	9.2	6.6	11.9	14.2	8.9	19.4	6.0	3.3	8.7
Capital (NS6)	4.3	3.0	5.6	5.9	3.4	8.4	3.4	2.0	4.9
New Brunswick	5.5	4.6	6.4	8.2	6.4	10	3.9	2.9	4.9
Moncton (NB1)	5.0	3.3	6.7	4.9	2.2	7.6	5.0	2.8	7.1
Sussex/Saint John (NB2)	5.7	3.8	7.6	9.3	5.5	13.2	3.6	1.5	5.7
Fredericton (NB3)	5.7	3.6	7.7	8.3	4.3	12.3	4.3	2.0	6.7
Edmundston (NB4)	7.2	3.3	11.2	12.3	4.2	20.3	3.8	--	7.5
Campbellton (NB5)	3.6	0.1	7.0	--	--	--	--	--	--
Bathurst (NB6)	6.6	3.6	9.6	12.0	5.7	18.3	2.9	0.3	5.5
Miramichi (NB7)	4.3	1.1	7.5	7.4	0.8	14.1	--	--	--

Table 160. All other respiratory disease deaths (ICD-9 460-519, excluding 480-487, 490-493), crude count and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts

	Both sexes			Male			Female		
	Crude count	95% CI		Crude count	95% CI		Crude count	95% CI	
		low	high		low	high		low	high
Canada	9,768	9,656	9,880	5,691	5,606	5,776	4,077	4,005	4,150
Newfoundland and Labrador	170	156	185	111	100	123	59	51	68
St. John's (NF1)	55	47	64	32	26	39	24	19	30
Eastern (NF2)	42	35	49	27	22	33	15	11	19
Central (NF3)	34	28	41	27	21	33	7	5	11
Western (NF4)	30	24	36	20	15	25	10	7	14
Grenfell (NF5)	3	2	6	2	1	5	1	--	3
Labrador (NF6)	6	4	9	4	2	6	2	1	5
Prince Edward Island	46	39	55	26	21	33	20	15	26
PEI/Urban	26	20	32	12	9	16	14	10	18
PEI/Rural	21	16	26	14	11	19	6	4	10
Nova Scotia	407	385	430	244	226	262	163	149	178
South-SW (NS1)	48	40	56	30	24	37	18	13	23
Valley (NS2)	36	30	44	23	18	29	13	9	18
Colch-Cumb-E.Hants (NS3)	56	48	65	30	24	36	26	21	32
Pictou-GASHA (NS4)	48	40	56	30	24	37	18	13	23
Cape Breton (NS5)	81	71	91	52	45	61	28	23	35
Capital (NS6)	139	126	153	78	69	89	61	52	70
New Brunswick	304	284	324	178	163	194	126	113	139
Moncton (NB1)	60	52	70	35	28	42	26	20	32
Sussex/Saint John (NB2)	88	78	99	47	40	55	41	34	48
Fredericton (NB3)	70	61	80	45	38	53	25	20	32
Edmundston (NB4)	23	18	29	14	10	19	9	6	13
Campbellton (NB5)	14	10	18	7	5	11	6	4	10
Bathurst (NB6)	30	24	36	20	15	26	10	7	14
Miramichi (NB7)	19	15	24	10	7	14	9	6	13

Table 161. All other respiratory disease deaths (ICD-9 460-519, excluding 480-487, 490-493), crude rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	32.9	32.5	33.3	38.7	38.2	39.3	27.2	26.7	27.7
Newfoundland and Labrador	30.3	27.8	33.0	39.7	35.6	44.2	21.0	18.1	24.2
St. John's (NF1)	29.7	25.4	34.4	34.6	28.1	42.0	24.9	19.6	31.3
Eastern (NF2)	33.4	28.0	39.7	43.6	34.8	53.9	23.4	17.2	31.1
Central (NF3)	30.0	24.6	36.3	46.7	37.2	57.8	13.1	8.4	19.4
Western (NF4)	32.0	25.9	39.2	42.5	32.6	54.4	21.6	14.8	30.4
Grenfell (NF5)	18.6	9.5	33.1	25.5	11.2	50.1	11.4	3.0	30.7
Labrador (NF6)	23.3	14.3	36.1	27.6	14.5	47.9	18.8	8.2	36.9
Prince Edward Island	34.0	28.7	40.0	39.2	31.2	48.6	29.0	22.3	37.0
PEI/Urban	40.4	32.1	50.2	38.6	27.4	52.8	42.1	30.6	56.6
PEI/Rural	28.4	22.0	36.2	39.7	29.1	52.9	17.3	10.7	26.5
Nova Scotia	43.7	41.3	46.2	53.3	49.6	57.3	34.4	31.5	37.6
South-SW (NS1)	37.4	31.6	43.9	47.6	38.5	58.2	27.4	20.7	35.5
Valley (NS2)	43.5	35.9	52.3	56.6	44.4	71.0	30.8	22.2	41.6
Colch-Cumb-E.Hants (NS3)	52.4	44.9	60.8	56.7	45.8	69.5	48.2	38.4	59.8
Pictou-GASHA (NS4)	47.7	40.3	56.0	60.8	49.1	74.3	34.9	26.4	45.3
Cape Breton (NS5)	56.6	49.8	64.0	75.5	64.4	88.0	38.6	31.0	47.5
Capital (NS6)	37.4	34.0	41.1	43.1	37.9	48.9	32.0	27.6	36.9
New Brunswick	40.3	37.8	43.0	47.6	43.7	51.8	33.1	29.9	36.6
Moncton (NB1)	33.0	28.4	38.1	38.3	31.4	46.2	27.8	22.1	34.5
Sussex/Saint John (NB2)	49.3	43.6	55.5	53.9	45.5	63.4	44.8	37.4	53.3
Fredericton (NB3)	42.5	37.1	48.6	54.5	45.8	64.3	30.6	24.3	38.1
Edmundston (NB4)	42.0	33.0	52.9	51.8	37.8	69.3	32.5	21.9	46.6
Campbellton (NB5)	41.5	30.1	55.7	45.1	29.0	67.0	38.0	23.5	58.1
Bathurst (NB6)	33.2	26.8	40.7	44.5	34.3	56.9	21.8	14.9	30.9
Miramichi (NB7)	38.0	29.1	48.9	39.6	27.2	55.8	36.4	24.5	52.1

Table 162. All other respiratory disease deaths (ICD-9 460-519, excluding 480-487, 490-493), age-standardized rate per 100,000 population and confidence interval, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	30.3	30.0	30.7	46.4	45.7	47.1	20.9	20.5	21.3
Newfoundland and Labrador	31.7	29.0	34.5	51.1	45.5	56.7	19.4	16.5	22.2
St. John's (NF1)	29.9	25.3	34.5	44.5	35.4	53.7	21.6	16.5	26.8
Eastern (NF2)	33.2	27.4	39.0	53.9	41.9	65.8	20.1	14.1	26.1
Central (NF3)	29.3	23.6	34.9	52.4	40.8	64.1	12.0	6.9	17.0
Western (NF4)	35.0	27.7	42.2	56.4	41.4	71.4	20.6	13.2	27.9
Grenfell (NF5)	26.0	9.6	42.4	42.8	8.7	76.9	--	--	--
Labrador (NF6)	69.1	34.6	103.6	86.0	30.9	141.2	53.7	10.8	96.6
Prince Edward Island	27.4	22.8	32.0	39.3	30.6	48.0	19.2	14.1	24.3
PEI/Urban	32.2	24.9	39.5	40.3	27.1	53.5	27.2	18.5	35.9
PEI/Rural	23.3	17.4	29.2	38.5	27.0	50.0	12.3	6.5	18.1
Nova Scotia	36.2	34.2	38.3	56.9	52.8	61.1	24.0	21.8	26.2
South-SW (NS1)	24.6	20.5	28.6	38.3	30.3	46.3	15.7	11.3	20.1
Valley (NS2)	32.9	26.7	39.2	55.2	42.3	68.1	20.3	13.7	26.9
Colch-Cumb-E.Hants (NS3)	37.9	32.1	43.7	52.1	41.3	62.9	27.9	21.6	34.3
Pictou-GASHA (NS4)	33.5	27.9	39.2	53.3	42.3	64.4	21.2	15.1	27.3
Cape Breton (NS5)	43.9	38.3	49.5	78.3	65.8	90.8	25.0	19.5	30.4
Capital (NS6)	39.7	35.9	43.5	61.2	53.3	69.2	28.2	24.1	32.4
New Brunswick	34.7	32.4	36.9	52.7	48.2	57.2	24.3	21.8	26.8
Moncton (NB1)	26.7	22.7	30.6	39.4	31.8	47.0	19.4	14.9	23.8
Sussex/Saint John (NB2)	39.8	35.0	44.7	58.6	48.8	68.5	30.8	25.2	36.5
Fredericton (NB3)	39.4	34.0	44.7	63.9	53.0	74.7	23.8	18.3	29.2
Edmundston (NB4)	37.6	28.8	46.5	60.6	42.1	79.1	24.5	15.2	33.9
Campbellton (NB5)	35.1	24.3	45.8	50.1	29.3	71.0	28.7	15.7	41.8
Bathurst (NB6)	31.3	24.8	37.9	53.1	39.7	66.5	17.1	10.8	23.4
Miramichi (NB7)	33.4	24.7	42.1	41.3	26.4	56.1	27.6	17.0	38.2

Table 163. Deaths due to medically treatable diseases - cervical cancer (ICD-9 180), age-standardized rate per 100,000 population and confidence interval, women aged 5 to 64, Canada, Atlantic Provinces, and Health Districts, 1996

	Female		
	Rate per 100,000	95% CI	
		low	high
Canada	1.8	1.7	1.9
Newfoundland and Labrador	2.9	1.6	4.2
St. John's (NF1)	2.6	0.5	4.7
Eastern (NF2)	F	F	F
Central (NF3)	2.8	0.1	5.5
Western (NF4)	5.3	1.0	9.5
Grenfell (NF5)	F	F	F
Labrador (NF6)	F	F	F
Prince Edward Island	F	F	F
PEI/Urban	F	F	F
PEI/Rural	F	F	F
Nova Scotia	2.9	2.0	3.9
South-SW (NS1)	2.9	0.3	5.4
Valley (NS2)	0.0	0.0	0.0
Colch-Cumb-E.Hants (NS3)	2.9	0.0	5.8
Pictou-GASHA (NS4)	F	F	F
Cape Breton (NS5)	5.4	2	8.8
Capital (NS6)	3	1.4	4.6
New Brunswick	1.8	1.0	2.7
Moncton (NB1)	F	F	F
Sussex/Saint John (NB2)	F	F	F
Fredericton (NB3)	3.8	1.2	6.5
Edmundston (NB4)	F	F	F
Campbellton (NB5)	F	F	F
Bathurst (NB6)	0.0	0.0	0.0
Miramichi (NB7)	F	F	F

Table 164. Deaths due to medically treatable diseases - bacterial infections (ICD-9 001-005, 020-041, 320, 382, 383, 390-392, 680-686, 711, 730), age-standardized rate per 100,000 population and confidence interval, population aged 5 to 64, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	0.8	0.7	0.9	0.8	0.8	0.9	0.8	0.7	0.8
Newfoundland and Labrador	0.6	0.2	1.0	0.8	0.2	1.5	F	F	F
St. John's (NF1)	F	F	F	F	F	F	0.0	0.0	0.0
Eastern (NF2)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Central (NF3)	F	F	F	F	F	F	F	F	F
Western (NF4)	F	F	F	0.0	0.0	0.0	F	F	F
Grenfell (NF5)	F	F	F	F	F	F	0.0	0.0	0.0
Labrador (NF6)	F	F	F	F	F	F	0.0	0.0	0.0
Prince Edward Island	F	F	F	F	F	F	0.0	0.0	0.0
PEI/Urban	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PEI/Rural	F	F	F	F	F	F	0.0	0.0	0.0
Nova Scotia	1.2	0.8	1.6	0.9	0.4	1.5	1.5	0.8	2.2
South-SW (NS1)	1.4	0.2	2.7	F	F	F	F	F	F
Valley (NS2)	1.8	0	3.6	F	F	F	F	F	F
Colch-Cumb-E.Hants (NS3)	F	F	F	F	F	F	F	F	F
Pictou-GASHA (NS4)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cape Breton (NS5)	1.3	0.2	2.4	F	F	F	2.1	0.0	4.2
Capital (NS6)	1.3	0.6	2.0	1.1	0.1	2.1	1.5	0.4	2.6
New Brunswick	0.7	0.4	1.1	0.7	0.1	1.2	0.8	0.2	1.4
Moncton (NB1)	0.9	0	1.8	F	F	F	F	F	F
Sussex/Saint John (NB2)	1.2	0.1	2.2	F	F	F	F	F	F
Fredericton (NB3)	F	F	F	F	F	F	F	F	F
Edmundston (NB4)	F	F	F	0.0	0.0	0.0	F	F	F
Campbellton (NB5)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Bathurst (NB6)	F	F	F	0.0	0.0	0.0	F	F	F
Miramichi (NB7)	F	F	F	0.0	0.0	0.0	F	F	F

Table 165. Deaths due to medically treatable diseases - pneumonia and unspecified bronchitis (ICD-9 481-486, 490), age-standardized rate per 100,000 population and confidence interval, population aged 5 to 49, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	0.8	0.8	0.9	1.0	0.9	1.1	0.7	0.6	0.8
Newfoundland and Labrador	0.6	0.2	1.0	1.0	0.2	1.7	F	F	F
St. John's (NF1)	F	F	F	F	F	F	F	F	F
Eastern (NF2)	F	F	F	F	F	F	0.0	0.0	0.0
Central (NF3)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Western (NF4)	F	F	F	F	F	F	0.0	0.0	0.0
Grenfell (NF5)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Labrador (NF6)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Prince Edward Island	F	F	F	F	F	F	0.0	0.0	0.0
PEI/Urban	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PEI/Rural	F	F	F	F	F	F	0.0	0.0	0.0
Nova Scotia	1.1	0.7	1.6	1.1	0.5	1.8	1.2	0.5	1.8
South-SW (NS1)	F	F	F	0.0	0.0	0.0	F	F	F
Valley (NS2)	F	F	F	0.0	0.0	0.0	F	F	F
Colch-Cumb-E.Hants (NS3)	F	F	F	0.0	0.0	0.0	F	F	F
Pictou-GASHA (NS4)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cape Breton (NS5)	F	F	F	0.0	0.0	0.0	F	F	F
Capital (NS6)	1.9	1.0	2.9	2.7	1.1	4.3	1.2	0.1	2.2
New Brunswick	0.4	0.1	0.6	F	F	F	0.5	0	0.9
Moncton (NB1)	F	F	F	F	F	F	0.0	0.0	0.0
Sussex/Saint John (NB2)	F	F	F	0.0	0.0	0.0	F	F	F
Fredericton (NB3)	F	F	F	F	F	F	F	F	F
Edmundston (NB4)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Campbellton (NB5)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Bathurst (NB6)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Miramichi (NB7)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Table 166. Deaths due to medically treatable diseases - hypertensive disease (ICD-9 401-405), age-standardized rate per 100,000 population and confidence interval, population aged 5 to 64, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	0.5	0.5	0.6	0.7	0.6	0.8	0.3	0.3	0.4
Newfoundland and Labrador	0.9	0.4	1.4	1.2	0.4	2.0	0.7	0.1	1.3
St. John's (NF1)	1.7	0.4	2.9	2.0	0.0	3.9	F	F	F
Eastern (NF2)	F	F	F	F	F	F	F	F	F
Central (NF3)	F	F	F	F	F	F	0.0	0.0	0.0
Western (NF4)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grenfell (NF5)	F	F	F	0.0	0.0	0.0	F	F	F
Labrador (NF6)	F	F	F	F	F	F	0.0	0.0	0.0
Prince Edward Island	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PEI/Urban	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
PEI/Rural	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
Nova Scotia	0.7	0.3	1.0	0.7	0.2	1.2	0.6	0.2	1.1
South-SW (NS1)	F	F	F	F	F	F	F	F	F
Valley (NS2)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Colch-Cumb-E.Hants (NS3)	F	F	F	0.0	0.0	0.0	F	F	F
Pictou-GASHA (NS4)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cape Breton (NS5)	1.6	0.3	2.9	2.2	0.0	4.3	F	F	F
Capital (NS6)	0.6	0.1	1.2	F	F	F	F	F	F
New Brunswick	0.3	0.1	0.6	0.5	0.1	1.0	F	F	F
Moncton (NB1)	F	F	F	F	F	F	0.0	0.0	0.0
Sussex/Saint John (NB2)	F	F	F	F	F	F	0.0	0.0	0.0
Fredericton (NB3)	F	F	F	F	F	F	F	F	F
Edmundston (NB4)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Campbellton (NB5)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Bathurst (NB6)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Miramichi (NB7)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Table 167. Thirty day acute myocardial infraction and 30 day stroke mortality 2002

	30 Day AMI Risk Adjusted†			30 Day Stroke Risk Adjusted†		
	Rate	95% CI		Rate	95% CI	
		low	high		low	high
Canada	12.6‡	19.2‡
Newfoundland and Labrador	24.4	22.5	26.3
St. John's (NF1)	20.6	17.1	24.2
Eastern (NF2)	34.9	31.1	38.8
Central (NF3)	24.4	20.5	28.3
Prince Edward Island	12.7	10.4	15.0	15.1	11.8	18.4
Nova Scotia	13.0	12.0	13.9	24.1	22.8	25.5
South-SW (NS1)	14.0	11.7	16.3	26.8	23.4	30.1
Colch-Cumb-E.Hants (NS3)	14.7	11.9	17.5	29.1	25.2	33.1
Cape Breton (NS5)	13.4	11.3	15.5	20.9	17.8	23.9
Capital (NS6)	11.4	9.8	13.0	21.3	18.7	23.8
New Brunswick	13.0	12.0	13.9	21.9	20.4	23.4
Moncton (NB1)	12.3	10.3	14.3	21.2	18.3	24.0
Sussex/Saint John (NB2)	12.3	10.2	14.3	19.4	16.2	22.6
Fredericton (NB3)	12.6	10.5	14.7	22.9	19.5	26.3

AMI 30-day in-hospital mortality data reported for Newfoundland in the 2001 report has since been found to be not comparable and therefore should be disregarded.

† 3-year pooled average: 1997-1999.

‡ The average rate includes only provinces/territories for which comparable data were available.

DISEASES

All data retrieved from:

<http://www4.statcan.ca/survey/invite3/proceed.cgi?loc=http://www.statcan.ca/english/freepub/82-221-XIE/00502/tables.htm>

Table 168. Arthritis/rheumatism, asthma, diabetes, and high blood pressure, Canada, Atlantic Provinces, and Health Districts, 2000/01

	Arthritis/rheumatism						Asthma						Diabetes						High Blood Pressure					
	Both sexes		Male		Female		Both sexes		Male		Female		Both sexes		Male		Female		Both sexes		Male		Female	
	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province
Canada																								
Newfoundland and Labrador	✓		×		✓		×		✓		✓		✓		×		×		✓		×		×	
St. John's (NF1)	×	×	✓	✓	✓	✓	×	×	✓	✓	×	×	✓	×	×	×	×	✓	✓	×	×	✓	×	✓
Eastern (NF2)	×	×	✓	✓	✓	✓	×	✓	×	✓	×	✓	✓	×	×	×	×	✓	✓	×	×	×	×	✓
Central (NF3)	✓	✓	×	×	×	×	×	×	✓	✓	✓	✓	✓	✓	×	×	×	×	✓	✓	×	×	×	×
Western (NF4)	✓	✓	×	×	×	×	×	×	✓	×	✓	✓	✓	×	-	-	×	×	✓	×	×	×	×	✓
Grenfell (NF5)	×	×	✓	✓	×	×	×	×	-	-	✓	✓	×	×	-	-	-	-	✓	×	×	✓	×	×
Labrador (NF6)	×	×	✓	✓	✓	✓	×	✓	✓	×	×	×	×	-	-	×	✓	✓	×	×	✓	✓	✓	✓
Prince Edward Island	✓		×	×	×		✓		×		s		✓		×		✓		✓		×		×	
PEI/Urban	✓	×	×	✓	×	✓	✓	×	×	s	s	✓	×	×	✓	✓	✓	✓	×	×	×	✓	✓	✓
PEI/Rural	✓	✓	×	×	×	×	×	×	✓	✓	s	s	✓	✓	×	×	×	×	✓	✓	×	✓	×	×
Nova Scotia	✓		×		×		✓		×		×		✓		×		×		✓		×		×	
South-SW (NS1)	✓	✓	×	×	×	×	✓	×	×	×	✓	✓	✓	✓	×	×	×	×	✓	✓	×	×	×	×
Valley (NS2)	✓	×	×	✓	×	✓	✓	s	×	✓	×	×	✓	×	✓	✓	×	s	✓	×	×	✓	×	✓
Colch-Cumb-E.Hants (NS3)	✓	✓	×	×	×	×	✓	×	✓	✓	×	×	✓	✓	×	×	×	×	✓	✓	×	×	×	×
Pictou-GASHA (NS4)	✓	×	×	✓	×	✓	×	×	✓	✓	×	×	✓	✓	×	×	×	×	✓	✓	×	✓	×	×
Cape Breton (NS5)	✓	✓	×	×	×	×	✓	✓	×	×	✓	✓	✓	✓	×	×	×	×	✓	✓	×	×	×	×
Capital (NS6)	✓	×	×	✓	×	✓	✓	✓	✓	✓	×	×	×	×	✓	✓	✓	✓	✓	×	×	✓	s	✓
New Brunswick	✓		×		×		✓		×		✓		✓		×		×		✓		×		×	×
Moncton (NB1)	✓	×	×	✓	×	✓	✓	×	×	×	×	✓	×	×	×	✓	✓	✓	✓	✓	×	×	×	×
Sussex/Saint John (NB2)	✓	✓	×	×	×	×	×	×	×	✓	✓	✓	✓	×	×	×	✓	✓	✓	✓	×	✓	×	×
Fredericton (NB3)	✓	×	×	✓	×	✓	✓	×	×	✓	✓	✓	✓	×	✓	×	×	×	×	×	✓	✓	✓	✓
Edmundston (NB4)	✓	✓	×	×	×	✓	✓	×	✓	✓	✓	✓	✓	×	×	×	s	✓	×	×	×	✓	✓	✓
Campbellton (NB5)	✓	✓	×	×	×	×	✓	✓	×	×	×	×	✓	✓	-	-	×	×	✓	✓	×	×	×	×
Bathurst (NB6)	✓	×	×	✓	×	✓	×	×	✓	✓	✓	✓	✓	×	-	-	×	×	✓	×	✓	✓	×	×
Miramichi (NB7)	✓	×	✓	✓	×	✓	×	×	✓	✓	-	-	✓	✓	-	-	×	×	✓	✓	×	×	×	×

× - lower percentage than the province or nation

✓ - higher percentage than the province or nation

s - same percentage as the province or nation

“-” - no data available for comparison

Table 169. Arthritis/rheumatism, by sex, household population aged 12 and over, Canada, Atlantic Provinces, and Health Districts, 2000/01

	With Arthritis/rheumatism						Without Arthritis/rheumatism					
	Both sexes		Male		Female		Both sexes		Male		Female	
	Number	%	Number	%	Number	%	Number	%	Number	%	Number	%
Canada	3,933,146	15.2	1,450,969	11.4	2,482,177	19.0	21,843,959	84.7	11,241,071	88.5	10,602,888	81.0
Newfoundland and Labrador	71,006	15.4	26,984	11.9	44,022	18.8	390,097	84.6	200,166	88.1	189,932	81.2
St. John's (NF1)	23,073	14.6	8,280	10.8	14,793	18.2	135,100	85.4	68,699	89.2	66,401	81.8
Eastern (NF2)	13,976	13.5	4,976 ^E	9.8 ^E	9,000	17.1	89,628	86.5	45,953	90.2	43,676	82.9
Central (NF3)	16,269	18.0	6,754	15.0	9,515	20.9	74,147	82.0	38,198	85.0	35,949	79.1
Western (NF4)	12,972	17.4	5,318	14.4	7,654	20.2	61,767	82.6	31,524	85.6	30,243	79.8
Grenfell (NF5)	1,975 ^E	14.1 ^E	644 ^E	9.1 ^E	1,330 ^E	19.3 ^E	12,045	85.9	6,469	90.9	5,576	80.7
Labrador (NF6)	2,742	13.6	1,012 ^E	9.8 ^E	1,730	17.6	17,409	86.4	9,322	90.2	8,086	82.4
Prince Edward Island	21,270	18.3	8,854	15.6	12,416	20.8	95,057	81.7	47,834	84.4	47,222	79.2
PEI/Urban	9,152	16.8	3,673	14.0	5,480	19.5	45,287	83.2	22,626	86.0	22,661	80.5
PEI/Rural	12,117	19.6	5,181	17.0	6,936	22.0	49,770	80.4	25,208	83.0	24,562	78.0
Nova Scotia	173,442	22.0	70,117	18.4	103,325	25.4	614,223	77.9	310,621	81.6	303,602	74.6
South-SW (NS1)	26,128	24.1	11,032	20.8	15,096	27.4	81,959	75.7	41,981	79.0	39,978	72.5
Valley (NS2)	13,815	19.6	5,427	16.0	8,388	23.0	56,582	80.3	28,542	84.0	28,040	76.8
Colch-Cumb-E.Hants (NS3)	22,360	25.1	9,786	22.6	12,573	27.6	66,577	74.9	33,575	77.4	33,002	72.4
Pictou-GASHA (NS4)	17,278	20.8	6,738	16.4	10,540	24.9	65,972	79.2	34,248	83.6	31,725	75.1
Cape Breton (NS5)	30,374	27.1	12,219	22.7	18,156	31.1	81,898	72.9	41,677	77.3	40,220	68.9
Capital (NS6)	63,487	19.6	24,914	16.0	38,572	22.8	261,235	80.4	130,598	84.0	130,637	77.2
New Brunswick	114,087	18.0	40,932	13.2	73,155	22.6	519,494	81.9	269,102	86.6	250,392	77.4
Moncton (NB1)	26,295	16.6	9,515	12.3	16,780	20.8	131,377	83.2	67,499	87.4	63,878	79.2
Sussex/Saint John (NB2)	31,877	21.3	10,091	14.0	21,786	28.3	117,463	78.7	62,206	86.0	55,257	71.7
Fredericton (NB3)	23,067	16.8	8,699	13.0	14,368	20.3	114,280	83.1	57,974	86.8	56,306	79.7
Edmundston (NB4)	8,575	18.6	3,757	16.6	4,819	20.5	37,557	81.4	18,917	83.4	18,640	79.5
Campbellton (NB5)	5,432	20.2	2,011	15.2	3,422	25.0	21,344	79.2	11,060	83.6	10,284	75.0
Bathurst (NB6)	12,195	16.2	4,594 ^E	12.2 ^E	7,601	20.3	63,019	83.8	33,112	87.8	29,906	79.7
Miramichi (NB7)	6,646	16.1	2,266	11.0	4,380	21.2	34,454	83.6	18,333	89.0	16,121	78.2

Table 170. Arthritis/rheumatism, by sex, household population aged 12 and over, Canada and Atlantic Provinces, 1994/95-1998/99

		With Arthritis/rheumatism						Without Arthritis/rheumatism					
		Both sexes		Male		Female		Both sexes		Male		Female	
		Number	%	Number	%	Number	%	Number	%	Number	%	Number	%
1994/95	Canada	3,036,663	12.7	1,122,387	9.5	1,914,277	15.7	20,884,821	87.2	10,644,025	90.4	10,240,796	84.2
	Newfoundland and Labrador	53,744	11.1	26,512	10.9	27,232	11.3	429,622	88.9	215,619	89.1	214,003	88.7
	Prince Edward Island	18,059	16.5	8,610	16.0	9,449	16.9	91,545	83.5	45,190	84.0	46,355	83.1
	Nova Scotia	143,682	18.8	52,201	14.1	91,481	23.2	614,926	80.5	314,830	85.0	300,096	76.2
	New Brunswick	92,039	14.7	33,882	11.0	58,157	18.2	534,266	85.3	273,514	89.0	260,752	81.8
1996/97	Canada	3,400,020	13.8	1,175,836	9.7	2,224,185	17.8	21,175,367	86.1	10,907,988	90.2	10,267,379	82.2
	Newfoundland and Labrador	65,688	13.7	28,813	12.1	36,875	15.4	411,887	86.2	209,378	87.9	202,509	84.5
	Prince Edward Island	19,918	17.7	9,388	17.0	10,530	18.3	92,738	82.3	45,838	83.0	46,900	81.7
	Nova Scotia	157,738	20.4	54,038	14.4	103,700	26.0	616,820	79.6	321,558	85.6	295,262	74.0
	New Brunswick	103,191	16.3	36,366	11.7	66,825	20.8	528,817	83.7	274,023	88.3	254,794	79.2
1998/99	Canada	3,801,350	15.3	1,376,046	11.2	2,425,304	19.2	21,097,211	84.7	10,877,316	88.7	10,219,895	80.7
	Newfoundland and Labrador	66,388	14.4	29,611	13.0	36,777	15.7	396,228	85.6	198,851	87.0	197,377	84.3
	Prince Edward Island	20,732	18.3	9,521	17.2	11,211	19.4	92,358	81.7	45,754	82.8	46,604	80.6
	Nova Scotia	160,562	20.7	55,092	14.8	105,469	26.3	613,287	79.3	318,161	85.2	295,127	73.7
	New Brunswick	104,993	16.7	31,729	10.3	73,264	22.9	521,622	83.2	275,633	89.7	245,989	77.0

Table 171. Diabetes, by sex, household population aged 12 and over, Canada, Atlantic Provinces, and Health Districts 2000/01

	With Diabetes						Without Diabetes					
	Both sexes		Male		Female		Both sexes		Male		Female	
	Number	%	Number	%	Number	%	Number	%	Number	%	Number	%
Canada	1,063,698	4.1	556,838	4.4	506,860	3.9	24,719,208	95.8	12,138,236	95.5	12,580,972	96.1
Newfoundland and Labrador	26,651	5.8	12,329	5.4	14,322	6.1	434,453	94.2	214,820	94.6	219,632	93.9
St. John's (NF1)	8,569	5.4	4,723 ^E	6.1 ^E	3,846 ^E	4.7 ^E	149,603	94.6	72,255	93.9	77,348	95.3
Eastern (NF2)	6,269	6.1	3,072 ^E	6.0 ^E	3,197 ^E	6.1 ^E	97,335	93.9	47,857	94.0	49,479	93.9
Central (NF3)	6,700	7.4	3,124 ^E	6.9 ^E	3,576 ^E	7.9 ^E	83,716	92.6	41,828	93.1	41,888	92.1
Western (NF4)	3,782	5.1	F	F	3,016 ^E	8.0 ^E	70,958	94.9	36,077	97.9	34,881	92.0
Grenfell (NF5)	529 ^E	3.8 ^E	F	F	F	F	13,491	96.2	6,859	96.4	6,632	96.0
Labrador (NF6)	802 ^E	4.0 ^E	F	F	412 ^E	4.2 ^E	19,349	96.0	9,944	96.2	9,404	95.8
Prince Edward Island	5,821	5.0	3,583	6.3	2,237	3.8	110,448	94.9	53,072	93.6	57,376	96.2
PEI/Urban	2,544	4.7	1,629 ^E	6.2 ^E	915 ^E	3.3 ^E	51,895	95.3	24,670	93.8	27,225	96.7
PEI/Rural	3,277	5.3	1,955 ^E	6.4 ^E	1,322 ^E	4.2 ^E	58,553	94.6	28,402	93.5	30,151	95.7
Nova Scotia	40,769	5.2	19,343	5.1	21,427	5.3	746,998	94.8	361,295	94.9	385,703	94.7
South-SW (NS1)	7,146	6.6	2,931	5.5	4,215	7.6	100,940	93.2	49,983	94.1	50,957	92.4
Valley (NS2)	3,272	4.6	1,337 ^E	3.9 ^E	1,935 ^E	5.3 ^E	67,228	95.4	32,631	96.1	34,596	94.7
Colch-Cumb-E.Hants (NS3)	6,246	7.0	2,860 ^E	6.6 ^E	3,386 ^E	7.4 ^E	82,691	93.0	40,502	93.4	42,189	92.6
Pictou-GASHA (NS4)	5,777 ^E	6.9 ^E	3,506 ^E	8.6 ^E	2,271 ^E	5.4 ^E	77,473	93.1	37,480	91.4	39,993	94.6
Cape Breton (NS5)	6,638	5.9	3,300 ^E	6.1 ^E	3,338 ^E	5.7 ^E	105,634	94.1	50,596	93.9	55,038	94.3
Capital (NS6)	11,690 ^E	3.6 ^E	5,409 ^E	3.5 ^E	6,281 ^E	3.7 ^E	313,032	96.4	150,103	96.5	162,928	96.3
New Brunswick	32,191	5.1	16,572	5.3	15,619	4.8	601,815	94.9	293,964	94.6	307,852	95.1
Moncton (NB1)	7,871	5.0	5,169	6.7	2,702 ^E	3.3 ^E	149,859	94.9	72,104	93.3	77,755	96.4
Sussex/Saint John (NB2)	6,679	4.5	3,874 ^E	5.4 ^E	2,805 ^E	3.6 ^E	142,661	95.5	68,423	94.6	74,238	96.4
Fredericton (NB3)	7,545	5.5	3,222 ^E	4.8 ^E	4,323 ^E	6.1 ^E	129,939	94.5	63,588	95.2	66,351	93.9
Edmundston (NB4)	2,445 ^E	5.3 ^E	1,313 ^E	5.8 ^E	1,132 ^E	4.8 ^E	43,688	94.7	21,361	94.2	22,327	95.2
Campbellton (NB5)	1,758 ^E	6.5 ^E	F	F	1,046 ^E	7.6 ^E	25,124	93.3	12,465	94.2	12,659	92.4
Bathurst (NB6)	3,390 ^E	4.5 ^E	F	F	2,002 ^E	5.3 ^E	71,823	95.5	36,319	96.3	35,504	94.7
Miramichi (NB7)	2,504 ^E	6.1 ^E	F	F	1,610 ^E	7.8 ^E	38,721	93.9	19,704	95.7	19,017	92.2

Table 172. Diabetes, by sex, household population aged 12 and over, Canada and Atlantic Provinces, 1994/95-1998/99

		With Diabetes						Without Diabetes					
		Both sexes		Male		Female		Both sexes		Male		Female	
		Number	%	Number	%	Number	%	Number	%	Number	%	Number	%
1994/95	Canada	722,491	3.0	359,458	3.1	363,032	3.0	23,198,994	96.9	11,406,954	96.8	11,792,040	96.9
	Newfoundland and Labrador	18,993 ^E	3.9 ^E	6,688 ^E	2.8 ^E	12,305 ^E	5.1 ^E	464,373	96.1	235,443	97.2	228,930	94.9
	Prince Edward Island	3,881 ^E	3.5 ^E	1,809 ^E	3.4 ^E	2,072 ^E	3.7 ^E	105,723	96.5	51,991	96.6	53,732	96.3
	Nova Scotia	27,474 ^E	3.6 ^E	14,877 ^E	4.0 ^E	12,597 ^E	3.2 ^E	731,133	95.7	352,154	95.1	378,979	96.3
	New Brunswick	17,773 ^E	2.8 ^E	5,428 ^E	1.8 ^E	12,344 ^E	3.9 ^E	608,532	97.2	301,968	98.2	306,565	96.1
1996/97	Canada	787,924	3.2	423,157	3.5	364,766	2.9	23,797,702	96.8	11,671,574	96.5	12,126,128	97.0
	Newfoundland and Labrador	21,919 ^E	4.6 ^E	F	F	12,309 ^E	5.1 ^E	455,657	95.4	228,582	96.0	227,075	94.8
	Prince Edward Island	3,628 ^E	3.2 ^E	1,447 ^E	2.6 ^E	2,181 ^E	3.8 ^E	109,028	96.8	53,779	97.4	55,249	96.2
	Nova Scotia	35,506 ^E	4.6 ^E	23,467 ^E	6.2 ^E	12,038 ^E	3.0 ^E	738,581	95.4	351,658	93.6	386,924	97.0
	New Brunswick	20,070 ^E	3.2 ^E	7,342 ^E	2.4 ^E	12,728 ^E	4.0 ^E	611,938	96.8	303,047	97.6	308,891	96.0
1998/99	Canada	865,908	3.5	481,833	3.9	384,075	3.0	24,041,852	96.5	11,777,254	96.1	12,264,598	96.9
	Newfoundland and Labrador	23,944	5.2	10,263 ^E	4.5 ^E	13,681 ^E	5.8 ^E	438,672	94.8	218,199	95.5	220,473	94.2
	Prince Edward Island	3,560 ^E	3.1 ^E	1,770 ^E	3.2 ^E	1,791 ^E	3.1 ^E	109,530	96.9	53,505	96.8	56,024	96.9
	Nova Scotia	34,108	4.4	22,189 ^E	5.9 ^E	11,919 ^E	3.0 ^E	739,741	95.6	351,064	94.1	388,677	97.0
	New Brunswick	20,708 ^E	3.3 ^E	6,792 ^E	2.2 ^E	13,917 ^E	4.4 ^E	606,189	96.7	300,570	97.8	305,618	95.6

Table 173. Asthma, by sex, household population aged 12 and over, Canada, Atlantic Provinces, and Health Districts 2000/01

	With Asthma						Without Asthma					
	Both sexes		Male		Female		Both sexes		Male		Female	
	Number	%	Number	%	Number	%	Number	%	Number	%	Number	%
Canada	2,170,748	8.4	878,920	6.9	1,291,828	9.9	23,615,991	91.5	11,818,983	93.0	11,797,008	90.1
Newfoundland and Labrador	30,477	6.6	11,973	5.3	18,504	7.9	430,582	93.4	215,131	94.7	215,450	92.1
St. John's (NF1)	10,267	6.5	3,714 ^E	4.8 ^E	6,552 ^E	8.1 ^E	147,906	93.5	73,264	95.2	74,642	91.9
Eastern (NF2)	7,806	7.5	2,852 ^E	5.6 ^E	4,954	9.4	95,798	92.5	48,076	94.4	47,722	90.6
Central (NF3)	5,735 ^E	6.3 ^E	2,262 ^E	5.0 ^E	3,473 ^E	7.6 ^E	84,636	93.6	42,646	94.9	41,990	92.4
Western (NF4)	4,496	6.0	2,307 ^E	6.3 ^E	2,189 ^E	5.8 ^E	70,243	94.0	34,535	93.7	35,708	94.2
Grenfell (NF5)	600 ^E	4.3 ^E	F	F	374 ^E	5.4 ^E	13,420	95.7	6,887	96.8	6,533	94.6
Labrador (NF6)	1,572 ^E	7.8 ^E	611 ^E	5.9 ^E	961 ^E	9.8 ^E	18,578	92.2	9,723	94.1	8,855	90.2
Prince Edward Island	9,877	8.5	3,945	7.0	5,932	9.9	106,432	91.5	52,742	93.0	53,690	90.0
PEI/Urban	4,722	8.7	1,923	7.3	2,798	9.9	49,718	91.3	24,375	92.7	25,343	90.1
PEI/Rural	5,156	8.3	2,022	6.7	3,134	9.9	56,715	91.6	28,368	93.3	28,347	90.0
Nova Scotia	72,022	9.1	28,524	7.5	43,498	10.7	715,879	90.9	352,248	92.5	363,631	89.3
South-SW (NS1)	9,763	9.0	5,798 ^E	10.9 ^E	3,965 ^E	7.2 ^E	98,529	91.0	47,322	89.1	51,207	92.8
Valley (NS2)	6,439	9.1	2,466 ^E	7.3 ^E	3,973 ^E	10.9 ^E	64,061	90.9	31,502	92.7	32,558	89.1
Colch-Cumb-E.Hants (NS3)	8,012	9.0	2,797 ^E	6.5 ^E	5,215	11.4	80,925	91.0	40,564	93.5	40,361	88.6
Pictou-GASHA (NS4)	6,784 ^E	8.1 ^E	2,060 ^E	5.0 ^E	4,724 ^E	11.2 ^E	76,396	91.8	38,855	94.8	37,541	88.8
Cape Breton (NS5)	10,476	9.3	5,077 ^E	9.4 ^E	5,399	9.2	101,796	90.7	48,819	90.6	52,977	90.8
Capital (NS6)	30,548	9.4	10,326 ^E	6.6 ^E	20,222	12.0	294,173	90.6	145,186	93.4	148,988	88.0
New Brunswick	54,447	8.6	22,994	7.4	31,453	9.7	579,644	91.4	287,541	92.6	292,103	90.2
Moncton (NB1)	16,414	10.4	6,116	7.9	10,299	12.8	141,517	89.6	71,157	92.1	70,360	87.2
Sussex/Saint John (NB2)	11,862	7.9	5,045 ^E	7.0 ^E	6,817	8.8	137,361	92.0	67,252	93.0	70,109	91.0
Fredericton (NB3)	12,058	8.8	5,597 ^E	8.4 ^E	6,461	9.1	125,427	91.2	61,214	91.6	64,213	90.9
Edmundston (NB4)	3,330 ^E	7.2 ^E	1,499 ^E	6.6 ^E	1,831 ^E	7.8 ^E	42,802	92.8	21,175	93.4	21,628	92.2
Campbellton (NB5)	2,780 ^E	10.3 ^E	1,225 ^E	9.3 ^E	1,555 ^E	11.3 ^E	24,102	89.5	11,951	90.3	12,150	88.7
Bathurst (NB6)	6,018	8.0	2,570 ^E	6.8 ^E	3,449 ^E	9.2 ^E	69,195	92.0	35,137	93.2	34,058	90.8
Miramichi (NB7)	1,984 ^E	4.8 ^E	943 ^E	4.6 ^E	F	F	39,240	95.2	19,656	95.4	19,585	94.9

Table 174. Asthma, by sex, household population aged 12 and over, Canada and Atlantic Provinces, 1994/95-1998/99

		With Asthma						Without Asthma					
		Both sexes		Male		Female		Both sexes		Male		Female	
		Number	%	Number	%	Number	%	Number	%	Number	%	Number	%
1994/95	Canada	1,955,472	7.2	977,483	7.3	977,989	7.1	25,050,943	92.5	12,363,390	92.4	12,687,553	92.6
	Newfoundland and Labrador	31,772	5.8	11,965	4.4	19,807 ^E	7.3 ^E	514,412	94.1	262,280	95.6	252,132	92.7
	Prince Edward Island	9,017	7.2	4,678	7.5	4,339	6.8	116,346	92.7	57,220	92.3	59,126	93.1
	Nova Scotia	69,438	8.0	38,884 ^E	9.2 ^E	30,554	6.9	788,574	91.3	379,079	90.0	409,494	92.6
	New Brunswick	46,984	6.7	19,442	5.6	27,542 ^E	7.7 ^E	658,518	93.3	328,314	94.3	330,204	92.2
1996/97	Canada	2,158,697	7.8	957,353	7.0	1,201,344	8.5	25,665,372	92.2	12,792,364	93.0	12,873,008	91.4
	Newfoundland and Labrador	31,622	5.9	16,069 ^E	5.9 ^E	15,553 ^E	5.8 ^E	508,580	94.1	254,797	94.1	253,784	94.1
	Prince Edward Island	9,808	7.6	4,474 ^E	7.0 ^E	5,334 ^E	8.2 ^E	119,311	92.4	59,442	93.0	59,869	91.8
	Nova Scotia	72,090	8.3	30,031 ^E	7.1 ^E	42,059	9.4	797,106	91.7	392,110	92.9	404,996	90.6
	New Brunswick	48,032 ^E	6.7 ^E	17,088 ^E	4.8 ^E	30,943 ^E	8.5 ^E	666,823	93.3	335,340	95.2	331,482	91.5
1998/99	Canada	2,389,085	8.5	1,096,815	7.9	1,292,270	9.1	25,690,359	91.5	12,786,267	92.1	12,904,092	90.8
	Newfoundland and Labrador	33,506	6.5	16,755 ^E	6.6 ^E	16,751 ^E	6.4 ^E	480,553	93.5	236,831	93.4	243,722	93.6
	Prince Edward Island	10,723	8.3	4,187 ^E	6.6 ^E	6,536	10.0	118,141	91.7	59,578	93.4	58,563	90.0
	Nova Scotia	88,448	10.2	32,677 ^E	7.6 ^E	55,770	12.6	781,642	89.8	395,526	92.3	386,117	87.4
	New Brunswick	57,923 ^E	8.2	21,822 ^E	6.3 ^E	36,101 ^E	10.2 ^E	646,355	91.8	326,861	93.7	319,493	89.8

Table 175. High blood pressure, by sex, household population aged 12 and over, Canada, Atlantic Provinces, and Health Districts 2000/01

	With High Blood Pressure						Without High Blood Pressure					
	Both sexes		Male		Female		Both sexes		Male		Female	
	Number	%	Number	%	Number	%	Number	%	Number	%	Number	%
Canada	3,257,198	12.6	1,443,291	11.4	1,813,907	13.9	22,497,665	87.2	11,233,033	88.4	11,264,632	86.0
Newfoundland and Labrador	70,876	15.4	32,345	14.2	38,531	16.5	389,928	84.6	194,734	85.7	195,194	83.4
St. John's (NF1)	22,042	13.9	9,974	13.0	12,068	14.9	135,984	86.0	67,005	87.0	68,979	85.0
Eastern (NF2)	16,304	15.7	7,765 ^E	15.2 ^E	8,539	16.2	87,300	84.3	43,163	84.8	44,137	83.8
Central (NF3)	16,972	18.8	7,052	15.7	9,920	21.8	73,444	81.2	37,901	84.3	35,543	78.2
Western (NF4)	10,759	14.4	5,262	14.3	5,497	14.5	63,898	85.5	31,581	85.7	32,318	85.3
Grenfell (NF5)	2,118 ^E	15.1 ^E	883 ^E	12.4 ^E	1,235 ^E	17.9 ^E	11,903	84.9	6,231	87.6	5,672	82.1
Labrador (NF6)	2,681	13.3	1,409 ^E	13.6 ^E	1,271	13.0	17,398	86.3	8,854	85.7	8,545	87.0
Prince Edward Island	16,313	14.0	7,523	13.3	8,790	14.7	99,889	85.9	49,093	86.6	50,796	85.2
PEI/Urban	7,065	13.0	3,527	13.4	3,538	12.6	47,301	86.9	22,699	86.3	24,602	87.4
PEI/Rural	9,248	14.9	3,996	13.1	5,252	16.7	52,587	85.0	26,394	86.9	26,194	83.2
Nova Scotia	127,845	16.2	55,100	14.5	72,745	17.9	659,503	83.7	325,330	85.4	334,173	82.1
South-SW (NS1)	20,307	18.8	8,003	15.1	12,304	22.3	87,985	81.2	45,116	84.9	42,868	77.7
Valley (NS2)	9,031	12.8	3,925	11.6	5,106	14.0	61,468	87.2	30,043	88.4	31,425	86.0
Colch-Cumb-E.Hants (NS3)	16,436	18.5	6,848	15.8	9,588	21.0	72,501	81.5	36,513	84.2	35,988	79.0
Pictou-GASHA (NS4)	13,704	16.5	5,684	13.9	8,020	19.0	69,547	83.5	35,302	86.1	34,244	81.0
Cape Breton (NS5)	24,377	21.7	10,199	18.9	14,179	24.3	87,827	78.2	43,630	81.0	44,197	75.7
Capital (NS6)	43,990	13.5	20,442	13.1	23,548	13.9	280,175	86.3	134,725	86.6	145,450	86.0
New Brunswick	91,865	14.5	38,280	12.3	53,584	16.6	541,426	85.4	271,522	87.4	269,904	83.4
Moncton (NB1)	24,336	15.4	10,547	13.6	13,789	17.1	132,987	84.2	66,303	85.8	66,685	82.7
Sussex/Saint John (NB2)	23,152	15.5	8,743	12.1	14,409	18.7	126,188	84.5	63,555	87.9	62,634	81.3
Fredericton (NB3)	15,779	11.5	6,904	10.3	8,875	12.6	121,539	88.4	59,740	89.4	61,799	87.4
Edmundston (NB4)	6,336	13.7	3,414	15.1	2,922 ^E	12.5 ^E	39,708	86.1	19,171	84.6	20,537	87.5
Campbellton (NB5)	5,045	18.7	2,086	15.8	2,958	21.6	21,837	81.1	11,090	83.8	10,747	78.4
Bathurst (NB6)	10,068	13.4	3,127	8.3	6,941	18.5	65,145	86.6	34,579	91.7	30,566	81.5
Miramichi (NB7)	7,149	17.3	3,459	16.8	3,689	17.9	34,022	82.5	17,084	82.9	16,937	82.1

Table 176. High blood pressure, by sex, household population aged 12 and over, Canada and Atlantic Provinces, 1994/95-1998/99

		With High Blood Pressure						Without High Blood Pressure					
		Both sexes		Male		Female		Both sexes		Male		Female	
		Number	%	Number	%	Number	%	Number	%	Number	%	Number	%
1994/95	Canada	2,077,788	8.7	861,818	7.3	1,215,970	10.0	21,843,696	91.2	10,904,594	92.6	10,939,103	89.9
	Newfoundland and Labrador	47,251	9.8	22,270	9.2	24,981	10.4	436,115	90.2	219,861	90.8	216,254	89.6
	Prince Edward Island	12,263	11.2	5,186	9.6	7,077	12.7	97,341	88.8	48,614	90.4	48,727	87.3
	Nova Scotia	99,546	13.0	38,430	10.4	61,116	15.5	659,061	86.3	328,600	88.7	330,461	84.0
	New Brunswick	62,748	10.0	24,096	7.8	38,652	12.1	563,557	90.0	283,300	92.2	280,257	87.9
1996/97	Canada	2,470,957	10.0	1,052,068	8.7	1,418,889	11.4	22,099,111	89.9	11,032,321	91.2	11,066,789	88.6
	Newfoundland and Labrador	50,430	10.6	24,287	10.2	26,143	10.9	427,146	89.4	213,904	89.8	213,241	89.0
	Prince Edward Island	13,522	12.0	5,209	9.4	8,313	14.5	99,134	88.0	50,017	90.6	49,117	85.5
	Nova Scotia	125,838	16.2	43,922	11.7	81,915	20.5	648,366	83.7	331,319	88.2	317,047	79.5
	New Brunswick	71,844	11.4	27,375E	8.8 ^E	44,468	13.8	560,164	88.6	283,014	91.2	277,151	86.2
1998/99	Canada	2,717,370	10.9	1,106,931	9.0	1,610,439	12.7	22,173,809	89.0	11,140,846	90.9	11,032,964	87.2
	Newfoundland and Labrador	51,452	11.1	23,302	10.2	28,150	12.0	411,164	88.9	205,160	89.8	206,004	88.0
	Prince Edward Island	15,654	13.8	6,536	11.8	9,118	15.8	97,249	86.0	48,617	88.0	48,632	84.1
	Nova Scotia	130,740	16.9	46,925	12.6	83,815	20.9	642,476	83.0	326,328	87.4	316,148	78.9
	New Brunswick	75,043	12.0	22,035E	7.2 ^E	53,008	16.6	551,854	88.0	285,327	92.8	266,527	83.4

Table 177. Age-standardized cancer (malignant neoplasms) incidence (ICD-9 140-208), Canada, Atlantic Provinces, and Health Districts, 1996

	All cancer						Lung cancer						Colorectal cancer						Prostate cancer		Female breast cancer			
	Both sexes		Male		Female		Both sexes		Male		Female		Both sexes		Male		Female		Canada	Province	Canada	Province		
	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province	Canada	Province		
Canada																								
Newfoundland and Labrador	✓		✓		✓		✓		✓		✓		×		×		×		✓		✓		✓	
St. John's (NF1)	✓	×	✓	×	✓	×	✓	×	✓	✓	✓	×	×	×	×	×	×	×	✓	✓	✓	✓	×	×
Eastern (NF2)	✓	×	×	×	✓	✓	✓	×	×	×	✓	×	×	×	×	×	×	×	×	✓	✓	✓	✓	✓
Central (NF3)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	×	×	×	×	×	×	✓	×	×	✓	✓	✓
Western (NF4)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	×	✓	✓	✓
Grenfell (NF5)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	-	-	×	✓	×	✓	-	-	✓	×	×	✓	×	×
Labrador (NF6)	✓	✓	✓	✓	✓	✓	✓	✓	-	-	-	-	✓	✓	×	✓	-	-	-	-	-	-	×	×
Prince Edward Island	×		×		×		×		×		×		×		×		×		×		×		✓	
Nova Scotia	×		×		×		×		×		×		×		×		×		×		×		×	
South-SW (NS1)	×	✓	✓	✓	×	✓	×	✓	×	✓	×	✓	×	✓	×	✓	×	✓	✓	✓	✓	✓	✓	✓
Valley (NS2)	✓	✓	✓	✓	✓	✓	×	✓	×	×	✓	✓	×	×	×	✓	×	×	×	✓	✓	✓	✓	✓
Colch-Cumb-E.Hants (NS3)	×	✓	×	✓	×	✓	×	×	×	×	✓	✓	×	×	×	✓	×	×	×	✓	✓	✓	✓	✓
Pictou-GASHA (NS4)	×	×	×	✓	×	×	×	×	×	✓	×	×	×	×	×	×	×	✓	×	✓	×	✓	×	×
Cape Breton (NS5)	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	✓	×	×	×	×	×	×
Capital (NS6)	×	✓	×	✓	×	✓	×	✓	×	✓	×	×	×	✓	×	✓	×	×	×	✓	×	✓	×	✓
New Brunswick	×		×		✓		×		×		✓		×		×		×		×		×		×	
Moncton (NB1)	×	✓	×	✓	✓	✓	✓	✓	×	✓	✓	✓	×	×	×	×	×	×	×	✓	✓	✓	✓	✓
Sussex/Saint John (NB2)	×	×	×	×	×	×	×	×	×	×	×	×	×	✓	✓	✓	×	✓	×	✓	×	✓	×	×
Fredericton (NB3)	×	✓	×	✓	✓	✓	×	✓	×	✓	✓	✓	×	×	✓	✓	×	×	×	×	×	×	×	×
Edmundston (NB4)	✓	✓	×	✓	×	✓	×	×	×	×	✓	✓	✓	✓	×	×	✓	✓	✓	✓	✓	✓	✓	✓
Campbellton (NB5)	×	✓	×	×	✓	✓	×	×	×	×	×	×	×	×	×	×	×	×	✓	✓	✓	✓	✓	✓
Bathurst (NB6)	×	✓	×	✓	✓	✓	×	✓	×	✓	✓	✓	✓	✓	×	✓	✓	✓	×	×	✓	✓	✓	✓
Miramichi (NB7)	×	✓	×	✓	✓	✓	✓	✓	✓	✓	×	×	✓	✓	✓	✓	×	×	×	✓	✓	✓	✓	✓

× - less favourable rate than the province or nation ✓ - more favourable rate than the province or nation "-" - no data available for comparison

Table 178. All cancer incidence (malignant neoplasms) (ICD-9 140-208), age-standardized rate per 100,000 population, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	384.5	383.3	385.8	453.6	451.5	455.7	338.5	336.9	340.1
Newfoundland and Labrador	347.0	338.0	356.0	402.0	387.6	416.4	306.4	294.7	318.1
St. John's (NF1)	366.6	350.4	382.8	421.0	394.8	447.3	333.4	312.4	354.4
Eastern (NF2)	371.2	351.5	390.9	455.6	423.3	487.9	305.3	280.8	329.7
Central (NF3)	340.2	321.1	359.4	383.0	353.6	412.3	303.6	278.3	329.0
Western (NF4)	292.2	272.0	312.4	332.6	300.8	364.5	263.3	236.7	289.9
Grenfell (NF5)	269.9	222.1	317.6	298.4	224.7	372.1	248.3	184.5	312.0
Labrador (NF6)	327.7	270.1	385.3	387.5	295.3	479.8	265.5	195.4	335.5
Prince Edward Island	406.6	387.7	425.5	491.9	461.0	522.7	347.2	323.0	371.3
Nova Scotia	410.6	403.4	417.7	489.8	478.0	501.6	358.6	349.4	367.8
South-SW (NS1)	391.0	373.3	408.6	451.4	423.7	479.1	346.3	323.2	369.5
Valley (NS2)	374.8	352.3	397.3	447.9	411.6	484.3	318.8	290.2	347.3
Colch-Cumb-E.Hants (NS3)	404.5	384.2	424.9	479.2	446.4	512.0	354.8	328.4	381.2
Pictou-GASHA (NS4)	412.9	391.6	434.2	488.3	454.2	522.4	364.3	336.5	392.2
Cape Breton (NS5)	445.6	427.2	464.0	568.4	536.7	600.1	371.8	348.7	394.9
Capital (NS6)	401.7	389.6	413.9	474.8	454.3	495.2	357.0	341.7	372.4
New Brunswick	406.3	398.2	414.3	500.7	487.4	514.1	338.2	328.2	348.2
Moncton (NB1)	404.5	388.5	420.4	497.7	471.3	524.2	336.7	316.6	356.7
Sussex/Saint John (NB2)	431.6	414.7	448.5	522.2	494.1	550.2	369.3	348.0	390.6
Fredericton (NB3)	386.6	369.5	403.7	470.0	441.6	498.3	325.6	304.2	347.0
Edmundston (NB4)	346.9	319.4	374.5	467.7	419.1	516.4	264.2	231.5	296.9
Campbellton (NB5)	387.9	351.1	424.7	517.8	453.6	581.9	295.1	251.2	339.0
Bathurst (NB6)	394.3	370.7	417.9	480.8	442.4	519.2	324.7	295.1	354.3
Miramichi (NB7)	387.6	357.1	418.1	468.0	418.9	517.1	327.7	288.5	366.8

Table 179. All cancer incidence (malignant neoplasms) (ICD-9 140-208), age-standardized rate per 100,000 population, by sex, Canada and Atlantic Provinces, 1976-2001

	Canada			Newfoundland and Labrador			Prince Edward Island			Nova Scotia			New Brunswick		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
1976	325.3	371.9	294.9	308.7	382.7	245.0	278.9	271.8	288.1	334.5	384.3	299.1	305.4	353.7	270.2
1977	339.8	391.4	306.0	327.5	392.1	274.1	281.8	295.7	275.2	336.7	376.2	311.2	321.6	367.3	290.2
1978	358.0	417.2	319.3	339.9	402.3	286.8	254.2	261.0	252.2	330.8	364.0	309.8	335.4	399.1	288.2
1979	351.6	409.8	313.8	319.7	376.6	272.2	313.2	341.9	297.9	338.5	385.3	307.6	325.1	381.8	284.0
1980	345.0	406.1	305.5	298.9	361.1	245.1	363.4	432.4	313.3	341.0	383.2	313.1	327.8	381.3	290.1
1981	372.5	442.1	328.1	299.1	355.2	254.4	368.0	434.6	322.9	323.0	374.9	286.2	327.2	395.0	280.0
1982	368.1	440.7	321.0	313.5	393.2	245.9	379.0	429.7	342.9	358.7	416.9	319.5	334.3	398.8	288.2
1983	377.6	448.4	332.8	324.8	371.2	289.4	354.9	398.7	322.2	349.2	414.1	301.8	335.9	409.6	285.2
1984	376.8	450.0	329.5	327.5	387.1	280.0	368.0	445.5	301.9	379.3	449.8	330.4	368.6	444.8	317.0
1985	379.5	449.8	335.5	325.2	388.1	276.3	354.3	403.9	319.2	378.6	440.4	337.9	367.0	451.7	309.1
1986	374.4	451.9	324.9	334.0	413.5	274.1	290.2	348.7	245.0	393.5	477.7	336.6	351.3	434.8	292.4
1987	379.6	456.3	330.7	332.1	393.5	283.8	368.3	466.5	294.0	396.3	472.7	342.6	386.0	475.0	326.6
1988	383.6	458.4	336.0	344.1	409.3	292.8	361.5	416.5	319.4	396.0	467.7	349.0	374.9	464.2	314.1
1989	377.4	451.5	330.0	343.6	398.4	303.4	400.3	483.0	334.5	407.9	474.2	365.3	371.2	462.9	306.3
1990	381.9	457.6	333.2	339.0	394.2	297.2	404.6	466.8	367.6	420.3	493.6	371.2	382.4	467.1	325.7
1991	388.9	469.0	337.1	330.3	403.2	274.8	378.7	415.1	352.6	418.8	503.7	362.4	380.8	459.4	327.6
1992	397.9	483.1	341.9	331.2	384.2	292.9	424.7	497.5	369.8	418.9	513.0	357.0	408.1	516.7	331.3
1993	404.7	497.1	341.8	352.0	433.7	289.3	414.0	537.6	318.2	444.4	557.0	365.1	427.2	545.3	345.9
1994	398.0	482.8	340.1	330.8	396.8	277.3	442.7	531.5	385.1	428.3	519.9	366.5	428.7	550.0	343.0
1995	386.4	458.6	338.4	361.1	421.3	316.5	412.1	508.5	343.1	415.4	498.1	361.0	401.0	503.3	325.9
1996	381.5	449.0	336.4	342.1	392.5	304.7	394.7	461.3	350.0	405.8	480.6	356.0	402.5	495.7	337.5
1997	384.8	451.8	340.0	338.5	392.6	298.8	412.7	501.8	349.8	409.6	488.8	358.3	414.1	501.9	349.6
1998	..	445.8	345.3	..	379.1	294.3	..	485.3	409.7	..	521.7	387.8	..	505.2	355.7
1999	..	444.9	344.0	..	344.3	243.4	..	493.3	336.2	..	516.7	371.6	..	488.6	351.2
2000	..	443.9	345.1	..	358.4	281.8	..	462.5	356.1	..	509.3	374.8	..	505.9	357.3
2001	..	443.0	346.2	..	354.1	281.0	..	466.3	356.1	..	511.7	376.8	..	509.6	361.0

Note: 1998 to 2001 figures are forecasts

Table 180. Lung cancer incidence (ICD-9 162), age-standardized rate per 100,000 population, by sex, Canada, Atlantic Provinces, and Health Districts 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	59.0	58.5	59.5	82.2	81.3	83.1	41.6	41.0	42.2
Newfoundland and Labrador	45.2	42.0	48.5	69.9	63.9	75.8	23.4	20.1	26.7
St. John's (NF1)	47.9	42.0	53.9	69.3	58.8	79.9	30.2	23.8	36.7
Eastern (NF2)	57.4	49.5	65.2	94.6	79.8	109.4	25.2	18.1	32.4
Central (NF3)	33.5	27.4	39.5	55.6	44.5	66.7	12.0	6.8	17.2
Western (NF4)	42.1	34.5	49.7	61.9	48.4	75.3	24.3	16.4	32.3
Grenfell (NF5)	31.2	14.8	47.6	57.3	26.0	88.7	x	x	x
Labrador (NF6)	32.6	15.5	49.8	x	x	x	x	x	x
Prince Edward Island	66.7	59.0	74.4	84.4	71.5	97.2	54.1	44.5	63.7
Nova Scotia	67.0	64.1	69.9	92.4	87.2	97.5	48.6	45.2	51.9
South-SW (NS1)	61.0	54.1	67.9	84.2	72.4	96.1	42.8	34.7	50.9
Valley (NS2)	63.5	54.2	72.8	93.3	76.8	109.8	39.9	29.6	50.3
Colch-Cumb-E.Hants (NS3)	67.1	58.8	75.3	99.2	84.3	114.1	40.5	31.8	49.2
Pictou-GASHA (NS4)	68.4	59.7	77.0	91.3	76.6	106.0	51.5	41.0	61.9
Cape Breton (NS5)	76.5	68.9	84.1	112.0	98.0	126.0	52.3	43.7	60.9
Capital (NS6)	64.5	59.6	69.5	82.9	74.2	91.5	52.8	46.8	58.8
New Brunswick	65.9	62.6	69.1	98.2	92.3	104.2	41.2	37.6	44.7
Moncton (NB1)	57.7	51.7	63.7	82.9	72.1	93.7	38.5	31.8	45.2
Sussex/Saint John (NB2)	73.8	66.9	80.8	105.2	92.6	117.8	52.1	44.1	60.2
Fredericton (NB3)	59.2	52.5	66.0	90.9	78.4	103.3	34.2	27.1	41.3
Edmundston (NB4)	76.0	63.1	88.8	138.1	111.6	164.6	30.4	19.5	41.4
Campbellton (NB5)	76.1	59.8	92.3	114.7	84.6	144.9	46.0	28.8	63.2
Bathurst (NB6)	61.8	52.2	71.3	97.1	79.6	114.5	30.6	21.3	39.8
Miramichi (NB7)	58.5	46.7	70.4	75.8	55.9	95.7	46.1	31.5	60.8

Table 181. Lung cancer incidence (ICD-9 162), age-standardized rate per 100,000 population, by sex, Canada and Atlantic Provinces, 1976-2001

	Canada			Newfoundland and Labrador			Prince Edward Island			Nova Scotia			New Brunswick		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
1976	43.3	75.7	16.3	33.5	57.1	9.9	31.4	50.5	13.3	46.5	81.7	15.7	41.5	71.6	14.8
1977	45.4	78.6	17.9	36.3	69.3	4.3	32.7	56.2	12.1	53.5	91.6	21.7	39.1	69.4	13.6
1978	49.3	85.1	20.1	40.3	67.3	14.6	25.6	41.4	11.8	51.5	87.4	20.7	47.5	83.8	17.1
1979	48.8	83.9	20.3	41.0	76.3	8.7	32.0	56.4	11.0	49.1	80.4	23.1	45.5	80.9	16.1
1980	49.0	83.2	21.7	36.9	66.1	8.9	43.0	70.0	19.1	55.2	87.7	28.2	44.5	72.2	20.7
1981	53.7	91.2	24.3	38.0	68.3	9.7	44.5	76.3	17.9	52.5	92.5	19.9	52.7	93.3	19.0
1982	55.3	92.6	25.9	44.7	81.9	9.8	48.1	87.9	13.4	54.7	87.5	28.0	56.1	94.2	24.9
1983	57.6	95.2	28.3	43.7	77.3	12.8	52.0	94.6	14.9	63.8	105.0	30.5	52.8	91.8	21.5
1984	59.1	97.1	29.6	44.3	77.4	13.9	66.0	117.1	22.5	65.6	107.4	32.5	62.9	109.6	24.7
1985	58.0	93.2	30.9	43.5	75.7	14.5	46.8	78.3	21.2	62.7	100.1	33.4	61.5	104.4	26.4
1986	59.6	96.4	31.6	49.0	88.0	13.5	52.3	83.9	24.9	64.1	104.4	33.0	57.2	97.4	25.3
1987	60.0	95.0	33.2	49.5	85.7	17.2	60.8	112.4	18.4	61.9	98.3	33.3	65.5	110.1	29.9
1988	61.1	95.5	34.8	57.1	100.9	16.6	45.0	69.9	23.5	67.5	103.0	40.5	61.3	102.3	29.3
1989	60.4	93.6	35.0	44.7	74.5	17.0	68.4	96.0	46.8	68.2	101.6	42.5	62.9	101.9	32.3
1990	60.9	92.7	36.5	45.1	73.5	20.8	65.5	77.9	56.2	67.3	103.8	39.5	66.9	108.2	34.5
1991	60.6	90.7	37.7	40.0	68.1	15.0	71.2	94.1	53.5	73.4	113.4	42.8	64.3	98.2	37.2
1992	61.4	90.3	39.6	42.4	68.1	20.0	59.1	75.5	47.3	70.0	105.5	43.4	69.4	104.8	42.1
1993	62.7	91.9	40.6	40.3	64.3	19.1	61.3	93.7	36.6	76.4	112.8	47.8	66.3	103.3	38.5
1994	60.2	87.3	39.8	39.2	63.9	16.9	78.9	115.8	50.5	68.1	99.0	44.7	68.6	107.0	38.5
1995	59.7	84.8	40.8	47.9	74.0	24.0	69.1	78.0	64.4	71.8	106.1	46.8	65.6	97.9	40.7
1996	59.2	82.3	42.0	44.6	69.7	22.3	68.4	85.1	54.9	62.3	84.8	45.2	66.9	100.2	42.1
1997	57.9	79.4	41.9	43.3	65.9	23.7	62.6	88.7	43.8	66.8	86.1	53.5	64.9	96.4	40.6
1998	..	79.3	42.9	..	58.0	25.6	..	102.9	54.2	..	101.5	49.6	..	95.7	49.7
1999	..	78.5	44.6	..	46.5	19.6	..	76.4	62.6	..	104.7	51.3	..	102.5	48.3
2000	..	76.9	45.5	..	51.1	23.8	..	87.6	56.0	..	96.2	53.7	..	98.6	49.5
2001	..	75.4	46.4	..	48.4	24.4	..	87.7	56.0	..	95.6	55.1	..	98.2	51.1

Note: 1998 to 2001 figures are forecasts

Table 182. Colorectal cancer incidence (ICD-9 153-154), age-standardized rate per 100,000 population, by sex, Canada, Atlantic Provinces, and Health Districts, 1996

	Both sexes			Male			Female		
	Rate per 100,000	95% CI		Rate per 100,000	95% CI		Rate per 100,000	95% CI	
		low	high		low	high		low	high
Canada	49.6	49.2	50.1	60.3	59.5	61.0	41.2	40.6	41.7
Newfoundland and Labrador	58.2	54.5	62.0	69.9	63.9	75.9	48.6	43.9	53.2
St. John's (NF1)	59.2	52.7	65.7	76.5	65.3	87.7	46.2	38.4	54.0
Eastern (NF2)	67.0	58.6	75.4	77.1	63.7	90.4	59.1	48.3	69.8
Central (NF3)	65.0	56.6	73.4	73.6	60.7	86.4	57.2	46.2	68.2
Western (NF4)	38.0	30.6	45.3	42.6	31.2	54.1	34.0	24.5	43.6
Grenfell (NF5)	50.5	29.5	71.4	69.6	33.3	105.9	x	x	x
Labrador (NF6)	39.8	19.5	60.1	62.7	26.1	99.2	x	x	x
Prince Edward Island	58.4	51.4	65.4	64.8	53.6	75.9	53.9	44.8	63.0
Nova Scotia	56.2	53.5	58.8	67.1	62.7	71.4	48.1	44.8	51.3
South-SW (NS1)	53.1	46.7	59.5	63.6	53.2	73.9	44.5	36.5	52.5
Valley (NS2)	57.0	48.4	65.6	65.0	51.2	78.8	49.3	38.6	60.0
Colch-Cumb-E.Hants (NS3)	60.0	52.4	67.7	67.0	54.8	79.3	54.9	45	64.7
Pictou-GASHA (NS4)	57.4	49.7	65.0	75.4	62.2	88.6	43.8	34.6	53.0
Cape Breton (NS5)	57.8	51.3	64.3	76.7	65.0	88.5	45.3	37.7	53.0
Capital (NS6)	53.8	49.3	58.2	61.0	53.6	68.4	48.6	42.9	54.2
New Brunswick	51.6	48.8	54.4	60.6	55.9	65.3	44.1	40.6	47.7
Moncton (NB1)	54.2	48.4	60.0	65.0	55.4	74.6	45.2	38.1	52.3
Sussex/Saint John (NB2)	50.7	44.9	56.4	59.9	50.3	69.4	43.0	36.0	50.0
Fredericton (NB3)	53.4	47.0	59.7	59.1	48.9	69.2	48.3	40.2	56.4
Edmundston (NB4)	46.8	36.8	56.8	65.9	47.5	84.3	32.9	21.8	44.1
Campbellton (NB5)	62.1	47.4	76.8	72.5	48.3	96.6	55.3	36.5	74.1
Bathurst (NB6)	39.7	32.2	47.1	47.7	35.7	59.8	32.9	23.6	42.1
Miramichi (NB7)	48.4	37.7	59.2	51.1	35.0	67.2	44.8	30.7	58.9

Table 183. Colorectal cancer incidence (ICD-9 153-154), age-standardized rate per 100,000 population, by sex, Canada and Atlantic Provinces, 1976-2001

	Canada			Newfoundland and Labrador			Prince Edward Island			Nova Scotia			New Brunswick		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
1976	49.3	55.3	44.6	43.5	58.2	30.7	41.9	38.1	44.0	59.5	65.9	54.6	42.4	40.6	43.8
1977	51.2	56.4	47.2	55.0	62.6	49.1	50.5	42.6	57.0	53.5	51.4	56.0	52.9	61.0	46.4
1978	53.7	59.4	49.5	55.4	57.5	54.0	48.5	46.8	49.9	53.1	53.8	52.3	54.8	61.2	48.8
1979	53.1	58.5	49.1	48.0	47.0	48.1	54.2	61.4	48.6	60.7	64.2	58.4	51.3	52.5	49.8
1980	51.2	57.3	46.8	55.4	60.1	50.9	53.4	63.2	47.0	58.6	55.7	61.2	60.1	67.0	54.8
1981	53.7	61.6	47.8	55.7	59.7	52.0	55.6	58.2	53.4	52.4	59.3	47.1	51.1	55.5	47.9
1982	53.8	61.9	48.0	56.6	64.7	50.0	48.8	49.9	47.1	52.4	63.3	43.6	53.2	56.7	50.1
1983	55.2	63.0	49.4	68.1	70.4	67.1	53.7	56.7	50.5	50.7	55.6	46.7	52.0	60.4	45.7
1984	55.0	64.0	48.3	55.4	63.6	48.1	55.9	50.7	59.3	56.9	64.4	50.3	56.1	59.9	53.4
1985	56.4	65.4	49.8	59.3	63.6	54.8	58.9	67.9	50.6	62.2	71.4	55.5	59.1	68.6	52.6
1986	54.5	63.8	47.4	60.9	67.8	56.1	51.0	72.4	32.3	62.3	73.9	54.0	55.0	63.0	48.5
1987	54.2	64.0	46.9	60.8	66.5	55.4	60.6	79.0	44.6	62.6	75.4	52.6	54.1	63.1	48.2
1988	53.4	63.7	45.4	54.2	64.9	45.8	58.8	66.7	50.7	59.0	66.7	52.0	59.4	72.0	49.1
1989	52.3	62.1	44.7	56.5	62.0	51.4	40.0	41.3	39.1	58.2	64.8	52.8	53.5	62.1	46.9
1990	52.5	62.2	45.0	55.3	62.1	50.0	54.2	58.5	51.0	64.0	75.3	54.7	53.2	60.8	47.6
1991	51.7	62.3	43.5	64.3	73.0	57.3	46.3	48.6	43.1	59.8	74.0	49.4	49.2	57.7	42.6
1992	52.1	63.4	43.4	62.6	69.2	56.8	72.5	73.8	70.7	57.9	68.5	51.1	57.5	69.7	47.7
1993	51.3	61.3	43.6	63.1	80.1	49.9	61.4	65.7	56.4	56.8	64.9	50.0	53.4	66.0	42.6
1994	51.6	62.4	43.1	58.9	69.1	49.5	55.8	54.9	57.4	57.2	66.9	49.9	54.3	69.2	42.8
1995	50.3	60.8	42.0	66.2	79.1	55.8	58.4	68.0	50.4	56.9	70.8	46.2	50.7	62.9	40.7
1996	49.2	60.1	40.4	53.6	67.2	42.1	57.1	55.7	60.2	56.0	66.7	48.6	50.5	60.5	42.1
1997	49.3	59.6	41.1	55.1	63.5	48.1	59.7	70.2	51.3	55.4	63.4	49.6	53.5	58.3	49.5
1998	..	61.3	43.1	..	70.2	52.4	..	78.0	64.6	..	76.4	58.0	..	64.0	49.6
1999	..	59.9	40.5	..	74.3	48.5	..	70.8	42.8	..	69.3	48.3	..	62.9	48.6
2000	..	59.6	40.1	..	72.5	48.1	..	67.1	53.7	..	67.8	49.1	..	62.1	45.1
2001	..	59.3	39.6	..	73.0	47.7	..	67.7	53.7	..	67.6	48.9	..	62.0	45.0

Note: 1998 to 2001 figures are forecasts

Table 184. Prostate cancer incidence (ICD-9 185), age-standardized rate per 100,000 population, Canada, Atlantic Provinces, and Health Districts, 1996

	Rate per 100,000	95% CI	
		low	high
Canada	112.2	111.1	113.2
Newfoundland and Labrador	90.1	83.2	97.1
St. John's (NF1)	90.4	77.9	102.8
Eastern (NF2)	92.1	77.3	106.8
Central (NF3)	101.8	86.6	117.0
Western (NF4)	81.9	65.6	98.1
Grenfell (NF5)	x	x	x
Labrador (NF6)	x	x	x
Prince Edward Island	142.8	126.1	159.5
Nova Scotia	117.8	112.0	123.6
South-SW (NS1)	94.0	81.4	106.6
Valley (NS2)	85.3	69.5	101.2
Colch-Cumb-E.Hants (NS3)	117.5	101.4	133.6
Pictou-GASHA (NS4)	114.5	98.1	130.9
Cape Breton (NS5)	158.5	141.7	175.3
Capital (NS6)	117.7	107.3	128.0
New Brunswick	134.2	127.3	141.2
Moncton (NB1)	127.5	114.0	141.0
Sussex/Saint John (NB2)	147.0	132.2	161.9
Fredericton (NB3)	134.9	119.6	150.2
Edmundston (NB4)	113.2	89.1	137.3
Campbellton (NB5)	106.6	77.7	135.6
Bathurst (NB6)	136.0	115.3	156.7
Miramichi (NB7)	117.2	92.4	141.9

Table 185. Prostate cancer incidence (ICD-9 185), age-standardized rate per 100,000 population, Canada and Atlantic Provinces, 1976-2001

	Canada	Newfoundland and Labrador	Prince Edward Island	Nova Scotia	New Brunswick
1976	62.1	58.0	63.8	65.3	71.4
1977	67.9	53.9	73.6	57.9	71.2
1978	74.0	49.1	48.2	62.1	69.0
1979	72.0	47.8	66.4	60.2	63.4
1980	71.4	46.3	81.4	60.0	61.5
1981	78.5	41.5	54.5	47.5	65.6
1982	77.8	46.9	72.5	73.1	66.5
1983	79.6	55.6	56.3	61.6	77.0
1984	80.9	54.3	69.4	76.7	84.8
1985	85.0	59.7	75.2	77.3	79.1
1986	86.1	61.4	56.1	78.5	81.9
1987	89.6	56.1	83.4	85.8	94.4
1988	90.4	60.5	82.2	88.9	100.1
1989	91.8	67.8	100.2	87.4	99.5
1990	99.8	61.9	108.5	83.1	100.1
1991	112.3	75.8	103.2	104.4	120.2
1992	125.3	67.5	147.0	119.1	134.7
1993	140.4	102.7	170.8	148.4	182.5
1994	129.4	80.8	201.4	145.4	163.9
1995	111.3	79.2	148.4	110.7	128.0
1996	109.7	95.1	112.4	112.4	130.5
1997	115.0	95.6	165.6	129.5	143.6
1998	113.7	100.2	127.3	131.4	150.2
1999	113.9	104.0	122.3	139.8	136.8
2000	116.0	95.0	136.6	137.6	144.7
2001	118.2	95.0	135.7	141.8	146.1

Table 186. Female breast cancer incidence (ICD-9 174), age-standardized rate per 100,000 population, Canada, Atlantic Provinces, and Health Districts, 1996

	Rate per 100,000	95% CI	
		low	high
Canada	99.9	99.0	100.7
Newfoundland and Labrador	88.6	82.3	94.8
St. John's (NF1)	103.3	91.6	115.1
Eastern (NF2)	78.7	66.4	91.1
Central (NF3)	83.3	70.2	96.4
Western (NF4)	73.2	59.3	87.1
Grenfell (NF5)	99.5	59.0	140.0
Labrador (NF6)	101.2	60.0	142.4
Prince Edward Island	98.2	85.2	111.2
Nova Scotia	103.4	98.5	108.4
South-SW (NS1)	99.3	86.8	111.8
Valley (NS2)	91.2	75.7	106.7
Colch-Cumb-E.Hants (NS3)	98.8	84.7	112.9
Pictou-GASHA (NS4)	112.8	97.0	128.6
Cape Breton (NS5)	104.4	92.1	116.7
Capital (NS6)	102.8	94.6	111.0
New Brunswick	100.3	94.8	105.8
Moncton (NB1)	95.1	84.4	105.9
Sussex/Saint John (NB2)	105.3	93.9	116.7
Fredericton (NB3)	110.7	98.1	123.2
Edmundston (NB4)	82.5	64.3	100.7
Campbellton (NB5)	67.9	46.7	89.1
Bathurst (NB6)	98.1	81.8	114.3
Miramichi (NB7)	92.6	71.6	113.7

Table 187. Female breast cancer incidence (ICD-9 174), age-standardized rate per 100,000 population, Canada and Atlantic Provinces, 1976-2001

	Canada	Newfoundland and Labrador	Prince Edward Island	Nova Scotia	New Brunswick
1976	84.6	69.3	85.9	81.8	74.0
1977	84.4	62.3	83.6	76.3	76.5
1978	86.1	72.3	74.9	77.1	76.9
1979	87.3	65.5	97.2	80.1	77.3
1980	83.3	62.5	88.8	83.4	72.6
1981	86.5	55.4	69.5	85.8	72.8
1982	86.0	55.3	97.3	93.8	81.8
1983	89.3	65.4	95.0	77.4	79.6
1984	90.3	76.5	100.8	88.0	87.4
1985	92.2	70.9	73.3	92.8	85.4
1986	88.6	64.2	55.2	90.5	78.1
1987	91.1	71.0	103.1	90.5	87.4
1988	97.8	75.7	104.8	94.1	89.0
1989	96.4	79.8	99.7	107.3	90.0
1990	96.0	80.9	107.1	100.4	91.0
1991	100.1	78.9	99.7	99.4	103.9
1992	102.0	86.2	97.8	99.8	94.6
1993	99.2	82.4	66.8	100.6	101.3
1994	98.9	81.3	119.6	105.3	101.5
1995	98.7	90.4	96.1	105.7	101.5
1996	98.5	90.7	95.3	99.5	99.6
1997	102.0	84.4	103.4	104.9	99.5
1998	102.6	86.2	136.8	112.1	103.0
1999	103.6	84.3	101.4	105.1	102.0
2000	104.4	91.4	106.9	109.5	107.3
2001	105.3	92.8	106.9	110.6	108.8

Table 188. Rate of positive HIV test reports (per 100,000 population), province/territory and year of test, Canada, 1999-2001

Province	1999	2000	2001
New Brunswick	0.53	1.32	1.19
Prince Edward Island and Nova Scotia	2.23	1.57	0.93
Newfoundland and Labrador	1.11	0.56	0.93
Canada	7.34	6.89	7.06

Source: Health Canada, Division of HIV/AIDS Epidemiology and Surveillance. HIV and AIDS in Canada. Surveillance report to December 31, 2001.
Centre for Infectious Disease Prevention and Control, Health Canada, 2002.

Table 189. Number of positive HIV test reports by province/territory and year of test, Canada, 1995-2001

Province	1995	1996	1997	1998	1999	2000	2001
New Brunswick	11	12	6	13	4	10	9
Prince Edward Island and Nova Scotia	37	37	34	32	24	17	10
Newfoundland and Labrador	7	10	7	14	6	3	5
Canada	2,364	2,113	1,913	1,761	1,672	1,626	1,646

Source: Health Canada, Division of HIV/AIDS Epidemiology and Surveillance. HIV and AIDS in Canada. Surveillance report to December 31, 2001.